

Study of compactness for compressible fluid models with a specific Korteweg tensor

Boris Haspot *

Abstract

This work is devoted to the global stability theory of solutions for a general isothermal model of capillary fluids derived by C. Rohde in [17], which can be used as a phase transition model.

This chapter is structured in the following way: first of all inspired by the result by P.-L. Lions in [15] on the Navier-Stokes compressible system we will show the global stability of weak solution for our system with isentropic pressure and next with general pressure. Next we will consider perturbations close to a stable equilibrium as in the case of strong solution.

1 Introduction

1.1 Presentation of the model

The correct mathematical description of liquid-vapor phase interfaces and their dynamical behavior in compressible fluid flow has a long history. We are concerned with compressible fluids endowed with internal capillarity. One of the first model which takes into consideration the variation of density on the interface between two phases, originates from the XIXth century work by Van der Waals and Korteweg [13]. It was actually derived in his modern form in the 1980s using the second gradient theory, see for instance [12, 18]. Korteweg suggests a modification of the Navier-Stokes system to account additionally for phase transition phenomena in introducing a term of capillarity. He assumed that the thickness of the interfaces was not null as in the *sharp interface approach*. This is called the *diffuse interface approach*.

Korteweg-type models are based on an extended version of nonequilibrium thermodynamics, which assumes that the energy of the fluid not only depends on standard variables but on the gradient of the density. In terms of the free energy, this principle takes the form of a generalized Gibbs relation, see [18].

In the present chapter, we follow a new approach introduced by Coquel, Rohde and their collaborators in [4]. They remark that the local diffuse interface approach requires more regular solution than in the original sharp interface approach. Indeed the interfaces are assumed of non zero thickness, so that the density vary continuously between the two

*Université Paris XII - Val de Marne 61, avenue du Général de Gaulle 94 010 CRETEIL Cedex
Téléphone : (33-1) 45 17 16 51 Télécopie : (33-1) 45 17 16 49 e-mail : haspot@univ-paris12.fr

interfaces, whereas in the sharp interface models, the interfaces represent zone of discontinuity for the density. Coquel, Rohde and their collaborators present an alternative model with a capillarity term which avoids spatial derivatives. The model reads:

$$(NSK) \quad \begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) - \mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u + \nabla(P(\rho)) = \kappa \rho \nabla D[\rho] \\ (\rho_{t=0}, u_{t=0}) = (\rho_0, u_0) \end{cases}$$

with:

$$\mu > 0 \quad \text{and} \quad \lambda + 2\mu > 0$$

where ρ denotes the density of the fluid and $u \in \mathbb{R}^N$ the velocity, μ and λ represent the viscosity coefficients, κ is a coefficient of capillarity, P is a general pressure function. We are particularly interested by Van der Waals type pressure:

$$P : (0, b) \rightarrow (0, +\infty) \\ P(\rho) = \frac{RT_* \rho}{b - \rho} - a \rho^2$$

where a, b, R, T_* are positive constants, R being the specific gas constant. For fixed values a, b we choose the constant reference temperature T_* so small as P to be monotone decreasing in some non-empty interval.

Further we impose the conditions:

$$u(t, x) \rightarrow 0, \quad \rho(t, x) \rightarrow 0 \quad \text{as} \quad |x| \rightarrow +\infty, \quad (1.1)$$

In last section, we consider also more general situations: monotone pressure, other boundary conditions than boundary condition 1.1, at infinity instead of 1.1:

$$u(t, x) \rightarrow u_\infty, \quad \rho(t, x) \rightarrow \rho_\infty \quad \text{as} \quad |x| \rightarrow +\infty, \quad (1.2)$$

where ρ_∞ is a given nonnegative constant.

The term $\kappa \rho \nabla D[\rho]$ corresponds to the capillarity which is supposed to model capillarity effects close to phase transitions in [13]. The classical Korteweg's capillarity term is $D[\rho] = \Delta \rho$.

Based on Korteweg's original ideas Coquel, Rohde and their collaborators in [4] and Rohde in [17] choose a nonlocal capillarity term D which penalizes rapid variations in the density field close from the interfaces. They introduce the following capillarity term:

$$D[\rho] = \phi * \rho - \rho$$

where ϕ is chosen so that:

$$\phi \in L^\infty(\mathbb{R}^N) \cap C^1(\mathbb{R}^N) \cap W^{1,1}(\mathbb{R}^N), \quad \int_{\mathbb{R}^N} \phi(x) dx = 1, \quad \phi \text{ even, and } \phi \geq 0.$$

This choice of capillarity term allows to get solution with jumps, i.e with sharp interfaces. Before tackling the global stability theory for the system (NSK) , let us derive formally the uniform bounds available on (ρ, u) .

1.2 Energy spaces

We assume to simplify that $P(\rho) = a\rho^\gamma$ with $\gamma \geq 1$. Let Π (free energy) be defined by:

$$\Pi(s) = s \left(\int_0^s \frac{P(z)}{z^2} dz \right), \quad (1.3)$$

so that $P(s) = s\Pi'(s) - \Pi(s)$, $\Pi'(\bar{\rho}) = 0$ and if we renormalize the mass equation:

$$\partial_t \Pi(\rho) + \operatorname{div}(u \Pi(\rho)) + P(\rho) \operatorname{div}(u) = 0 \quad \text{in } \mathcal{D}'((0, T) \times \mathbb{R}^N).$$

Notice that Π is convex. Multiplying the equation of momentum conservation by u and integrating by parts over $\mathbb{R}^N$, we obtain the following energy estimate:

$$\begin{aligned} \int_{\mathbb{R}^N} \left(\frac{1}{2} \rho |u|^2 + \Pi(\rho) + E_{global}[\rho(\cdot, t)] \right)(x) dx(t) + \int_0^t \int_{\mathbb{R}^N} (\mu D(u) : D(u) \\ + (\lambda + \mu) |\operatorname{div} u|^2) dx \leq \int_{\mathbb{R}^N} \left(\frac{|m_0|^2}{2\rho} + \Pi(\rho_0) + E_{global}[\rho_0] \right) dx, \end{aligned} \quad (1.4)$$

where we have:

$$E_{global}[\rho(\cdot, t)](x) = \frac{\kappa}{4} \int_{\mathbb{R}^N} \phi(x - y) (\rho(y, t) - \rho(x, t))^2 dy.$$

The only non-standard term is the energy term E_{global} which comes from the product of u with the capillarity term $\kappa \rho \nabla(\phi * \rho - \rho)$. Indeed we have:

$$\begin{aligned} \kappa \int_{\mathbb{R}^N} u(t, x) \rho(t, x) \cdot \nabla([\phi * \rho(t, \cdot)](x) - \rho(t, x)) dx \\ = -\kappa \int_{\mathbb{R}^N} \operatorname{div}(u(t, x) \rho(t, x)) ([\phi * \rho(t, \cdot)](x) - \rho(t, x)) dx, \\ = \kappa \int_{\mathbb{R}^N} \frac{\partial}{\partial t} \rho(t, x) ([\phi * \rho(t, \cdot)](x) - \rho(t, x)) dx, \\ = -\frac{d}{dt} \int_{\mathbb{R}^N} E_{global}[\rho(t, \cdot)](x) dx. \end{aligned}$$

To derive the last equality we use the relation:

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^N} E_{global}[\rho(t, \cdot)](x) dx &= \frac{\kappa}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \phi(x - y) (\rho(t, y) - \rho(t, x)) \frac{\partial}{\partial t} \rho(t, y) dy dx \\ &\quad + \frac{\kappa}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \phi(y - x) (\rho(t, x) - \rho(t, y)) \frac{\partial}{\partial t} \rho(t, x) dy dx, \\ &= \kappa \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \phi(x - y) (\rho(t, y) - \rho(t, x)) \frac{\partial}{\partial t} \rho(t, y) dy dx, \\ &= -\kappa \int_{\mathbb{R}^N} ([\phi * \rho(t, \cdot)](x) - \rho(t, x)) \frac{\partial}{\partial t} \rho(t, x) dx. \end{aligned}$$

where we just use integration by parts.

In the sequel we will note:

$$\mathcal{E}(\rho, \rho u)(t) = \int_{\mathbb{R}^N} \left(\frac{1}{2} \rho |u|^2 + \Pi(\rho) + E_{global}[\rho(\cdot, t)] \right)(x) dx(t), \quad (1.5)$$

We are interested to use the above inequality energy to determine the functional space we must work with.

So if we expand $E_{global}[\rho(\cdot, t)](x)$ we get:

$$E_{global}[\rho(t, \cdot)](x) = \frac{\kappa}{4}(\rho^2 + \phi * \rho^2 - 2\rho(\phi * \rho)).$$

Because by the mass equation we obtain that ρ is bounded in $L^\infty(0, T; L^1(\mathbb{R}^N))$ if we suppose that $\rho_0 \in L^1$ and we have supposed that $\phi \in L^\infty(\mathbb{R}^N)$, we obtain that $\rho(\phi * \rho)$ is bounded in $L^\infty(0, T; L^1(\mathbb{R}^N))$. So we get that $\rho^2 + \phi * \rho^2 \in L^\infty(0, T; L^1(\mathbb{R}^N))$ and as $\phi \geq 0$ and $\rho \geq 0$ we get a control of ρ in $L^\infty(0, T; L^2(\mathbb{R}^N))$ (a property which turns out to be important to taking advantage of the theory of renormalized solutions, indeed ρ in $L^\infty(0, T; L^2(\mathbb{R}^N))$ implies that $\rho \in L^2_{loc}(\mathbb{R}^+ \times \mathbb{R}^N)$ and we will can use the theorem of Diperna-Lions on renormalized solutions, see [14])

In view of (1.4), we can specify initial conditions on $\rho|_{t=0} = \rho_0$ and $\rho u|_{t=0} = m_0$ where we assume that:

- $\rho_0 \geq 0$ a.e in $\mathbb{R}^N$, $\rho_0 \in L^1(\mathbb{R}^N) \cap L^s(\mathbb{R}^N)$ with $s = \max(2, \gamma)$,
 - $m_0 = 0$ a.e on $\rho_0 = 0$,
 - $\frac{|m_0|^2}{\rho_0}$ (defined to be 0 on $\rho_0 = 0$) is in $L^1(\mathbb{R}^N)$.
- (1.6)

We deduce the following a priori bounds which give us the energy space in which we will work:

- $\rho \in L^\infty(0, T; L^1(\mathbb{R}^N) \cap L^s(\mathbb{R}^N))$,
- $\rho|u|^2 \in L^\infty(0, T; L^1(\mathbb{R}^N))$,
- $\nabla u \in L^2((0, T) \times \mathbb{R}^N)^N$.

We will use this uniform bound in our result of compactness. Let us emphasize at this point that the above a priori bounds do not provide any control on $\nabla \rho$ in contrast with the case of $D[\rho] = \Delta \rho$ (see in [6]).

1.3 Notion of weak solutions

We now explain what we mean by renormalized weak solutions, weak solutions, and bounded energy weak solution of problem (NSK).

Multiplying mass equation by $b'(\rho)$, we obtained the so-called renormalized equation (see [14]):

$$\frac{\partial}{\partial t} b(\rho) + \operatorname{div}(b(\rho)u) + (\rho b'(\rho) - b(\rho)) \operatorname{div} u = 0. \quad (1.7)$$

with:

$$b \in C^0([0, +\infty)) \cap C^1((0, +\infty)), \quad |b'(t)| \leq ct^{-\lambda_0}, \quad t \in (0, 1], \quad \lambda_0 < 1 \quad (1.8)$$

with growth conditions at infinity:

$$|b'(t)| \leq ct^{\lambda_1}, \quad t \geq 1, \quad \text{where } c > 0, \quad -1 < \lambda_1 < \frac{s}{2} - 1. \quad (1.9)$$

Definition 1.1 A couple (ρ, u) is called a renormalized weak solution of problem (NSK) if we have:

- Equation of mass holds in $\mathcal{D}'(\mathbb{R}^N)$.
- Equation (1.7) holds in $\mathcal{D}'(\mathbb{R}^N)$ for any function b belonging to (1.8) and (1.9).

Definition 1.2 Let the couple (ρ_0, u_0) satisfy;

- $\rho_0 \in L^1(\mathbb{R}^N)$, $\Pi(\rho_0) \in L^1(\mathbb{R}^N)$, $E_{global}[\rho_0] \in L^1(\mathbb{R}^N)$, $\rho_0 \geq 0$ a.e in $\mathbb{R}^N$.
- $\rho_0 u_0 \in (L^1(\mathbb{R}^N))^N$ such that $\rho_0 |u_0|^2 1_{\rho_0 > 0} \in L^1(\mathbb{R}^N)$
- and such that $\rho_0 u_0 = 0$ whenever $x \in \{\rho_0 = 0\}$,

where the quantity Π is defined in (1.3). We have the following definitions:

1. A couple (ρ, u) is called a weak solution of problem (NSK) on $\mathbb{R}$ if:

- (a) $\rho \in L^r(L^r(\mathbb{R}^N))$ for $s \leq r \leq +\infty$,
- (b) $P(\rho) \in L^\infty(L^1(\mathbb{R}^N))$, $\rho \geq 0$ a.e in $\mathbb{R} \times \mathbb{R}^N$,
- (c) $\nabla u \in L^2(L^2(\mathbb{R}^N))$, $\rho |u|^2 \in L^\infty(L^1(\mathbb{R}^N))$.
- (d) Mass equation holds in $\mathcal{D}'(\mathbb{R} \times \mathbb{R}^N)$.
- (e) Momentum equation holds in $\mathcal{D}'(\mathbb{R} \times \mathbb{R}^N)^N$.
- (f) $\lim_{t \rightarrow 0^+} \int_{\mathbb{R}^N} \rho(t) \varphi = \int_{\mathbb{R}^N} \rho_0 \varphi$, $\forall \varphi \in \mathcal{D}(\mathbb{R}^N)$,
- (g) $\lim_{t \rightarrow 0^+} \int_{\mathbb{R}^N} \rho u(t) \cdot \phi = \int_{\mathbb{R}^N} \rho_0 u_0 \cdot \phi$, $\forall \phi \in \mathcal{D}(\mathbb{R}^N)^N$.

2. A couple (ρ, u) is called a bounded energy weak solution of problem (NSK) if in addition to (1d), (1e), (1f), (1g) we have:

- The quantity $\mathcal{E}_0$ is finite and inequality (1.4) with $\mathcal{E}$ defined by (1.5) and with $\mathcal{E}_0$ in place of $\mathcal{E}(\rho(0), \rho u(0))$ holds a.e in $\mathbb{R}$.

1.4 Mathematical results

We wish to prove global stability results for (NSK) with $D[\rho] = \phi * \rho - \rho$ in functional spaces very close to energy spaces. In the non capillary case and $P(\rho) = a\rho^\gamma$, P.-L. Lions in [15] proved the global existence of weak solutions (ρ, u) to (NSK) with $\kappa = 0$ (which become the system of Navier-Stokes compressible isotherm) for $\gamma > \frac{N}{2}$ if $N \geq 4$, $\gamma \geq \frac{3N}{N+2}$ if $N = 2, 3$ and initial data (ρ_0, m_0) such that:

$$\rho_0, \rho_0^\gamma, \frac{|m_0|^2}{\rho_0} \in L^1(\mathbb{R}^N).$$

where we agree that $m_0 = 0$ on $\{x \in \mathbb{R}^N / \rho_0(x) = 0\}$. More precisely, he obtains the existence of global weak solutions (ρ, u) to (NSK) with $\kappa = 0$ such that for all $t \in (0, +\infty)$:

- $\rho \in L^\infty(0, T; L^\gamma(\mathbb{R}^N))$ and $\rho \in C([0, T], L^p(\mathbb{R}^N))$ if $1 \leq p < \gamma$,
- $\rho \in L^q((0, T) \times \mathbb{R}^N)$ for $q = \gamma - 1 + \frac{2\gamma}{N} > \gamma$.
- $\rho|u|^2 \in L^\infty(0, T; L^1(\mathbb{R}^N))$ and $Du \in L^2((0, T) \times \mathbb{R}^N)$.

Notice that the main difficulty for proving Lions' theorem consists in exhibiting strong compactness properties of the density ρ in $L^p_{loc}(\mathbb{R}^+ \times \mathbb{R}^N)$ spaces required to pass to the limit in the pressure term $P(\rho) = a\rho^\gamma$.

Let us mention that Feireisl in [8] generalized the result to $\gamma > \frac{N}{2}$ in establishing that we can obtain renormalized solution without imposing that $\rho \in L^2_{loc}(\mathbb{R}^+ \times \mathbb{R}^N)$ (what needed Lions in dimension $N = 2, 3$, that's why $\gamma - 1 + \frac{2\gamma}{N} \geq 2$), for this he introduces the concept of oscillation defect measure evaluating the loss of compactness. We refer to the book of Novotný and Straškraba for more details (see [16]).

Let us mention here that the existence of strong solution with $D[\rho] = \Delta\rho$ is known since the work by Hattori and Li in [9], [10] in the whole space $\mathbb{R}^N$. In [6], Danchin and Desjardins study the well-posedness of the problem for the isothermal case with constant coefficients in critical Besov spaces. We recall too the results from Rohde in [17] who obtains the existence and uniqueness in finite time for two-dimensional initial data in $H^4(\mathbb{R}^2) \times H^4(\mathbb{R}^2)$.

In the present chapter, we aim at showing the global stability of weak solutions in the energy spaces for the system (NSK). This work is composed of four parts, the first one concerns estimates on the density to get a gain of integrability on the density needed to pass to the weak limit in the term of pressure and of capillarity. The second part is the passage to the weak limit in the non-linear terms of the density and the velocity according to Lions' methods. The idea is to use renormalized solution to test the weak limit on convex test functions. In this part we will concentrate on the case of simple pressure of type $P(\rho) = a\rho^\gamma$. We get the following theorem where $(\rho_n, u_n)_{n \in \mathbb{N}}$ is a suit of global regular approximate solution for the problem (NSK).

Theorem 1.1 *Let $N \geq 2$. Let $\gamma > N/2$ if $N \geq 4$ and $\gamma \geq 1$ else.*

Let the couple (ρ_0^n, u_0^n) satisfy:

- ρ_0^n is uniformly bounded in $L^1(\mathbb{R}^N) \cap L^s(\mathbb{R}^N)$ with $s = \max(\gamma, 2)$ and $\rho_0^n \geq 0$ a.e in $\mathbb{R}^N$,
- $\frac{|\rho_0^n u_0^n|^2}{\rho_0^n}$ is uniformly bounded in $L^1(\mathbb{R}^N)$,
- and such that $\rho_0^n u_0^n = 0$ whenever $x \in \{\rho_0 = 0\}$.

In addition we suppose that ρ_0^n converges in $L^1(\mathbb{R}^N)$ to ρ_0 , then up to a subsequence (ρ_n, u_n) converges strongly to a weak solution (ρ, u) of the system (NSK) satisfying the initial condition (ρ_0, u_0) as in (1.6). Moreover we have the following convergence:

- $\rho_n \rightarrow_n \rho$ in $C([0, T], L^p(\mathbb{R}^N)) \cap L^r((0, T) \times K)$ for all $1 \leq p < s$, $1 \leq r < q$, with $q = s + \frac{N\gamma}{2} - 1$ if $N \geq 3$, where $K = \mathbb{R}^N$ except when $N \geq 4$ and $\gamma < \frac{N}{2}(1 + \frac{1}{N})$ where K is an arbitrary compact.
- $\rho_n \rightarrow_n \rho$ in $C([0, T], L^p(\mathbb{R}^2)) \cap L^r((0, T) \times K)$ for all $1 \leq p < s$, $1 \leq r < q$, with K an arbitrary compact in $\mathbb{R}^2$ if $N = 2$.

In addition we have:

- $\rho_n u_n \rightarrow \rho u$ in $L^p(0, T; L^r(\mathbb{R}^N))$ for all $1 \leq p < +\infty$ and $1 \leq r < \frac{2s}{s+1}$,
- $\rho_n(u_i)_n(u_j)_n \rightarrow \rho_n u_i u_j$ in $L^p(0, T; L^1(\Omega))$ for all $1 \leq p < +\infty$, $1 \leq i, j \leq N$ if $N \geq 3$, where $K = \mathbb{R}^N$ except when $N \geq 4$ and $\gamma < \frac{N}{2}(1 + \frac{1}{N})$ where K is an arbitrary compact.
- $\rho_n(u_i)_n(u_j)_n \rightarrow \rho_n u_i u_j$ in $L^p(0, T; L^1(\Omega))$ for all $1 \leq p < +\infty$, $1 \leq i, j \leq N$ with Ω an arbitrary bounded open set in $\mathbb{R}^2$ if $N = 2$.

In the third part we will focus on general pressure, and particular van der Waal's pressure. In the fourth part we concentrate on the case with initial data close to a constant $\bar{\rho}$, and we will work in Orlicz space, this case is the most adapted for the strong solution because it enables us to control the vacuum so that one can use the property of ellipticity of the momentum equation.

2 Existence of weak solution for a isentropic pressure law

2.1 A priori estimates on the density

In this part we are interested by getting a gain of integrability on the density and we consider the case where $P(\rho) = a\rho^\gamma$. This will enable us to pass to the weak limit in the pressure and the Korteweg terms. It is expressed by the following theorem:

Theorem 2.2 *Let $N \geq 2$ and $\gamma \geq 1$, with in addition $\gamma > \frac{N}{2}$ if $N \geq 4$. Let (ρ, u) be a regular solution of the system (NSK) with $\rho \geq 0$ and $\rho \in L^\infty(L^1 \cap L^{s+\varepsilon})$ where we define ε below. Then we have if $\gamma \geq \frac{N}{2}(1 + \frac{1}{N})$ for $N \geq 4$:*

$$\int_{(0,T) \times \mathbb{R}^N} (\rho^{\gamma+\varepsilon} + \rho^{2+\varepsilon}) dx dt \leq M \text{ for any } 0 < \varepsilon \leq \frac{2}{N}\gamma - 1 \text{ if } N \geq 4,$$

$$\text{and } 0 < \varepsilon \leq \frac{4}{N} - 1 \text{ if } N = 2, 3.$$

with M depending only on the initial conditions and on the time T .

If $\gamma < \frac{N}{2}(1 + \frac{1}{N})$ for $N \geq 4$, we have:

$$\int_{(0,T) \times K} (\rho^{\gamma+\varepsilon} + \rho^{2+\varepsilon}) dx dt \leq M' \text{ for any } 0 < \varepsilon \leq \frac{2}{N}\gamma - 1 \text{ if } N \geq 4,$$

for any arbitrary compact K with M' depending only on the initial conditions, on K and on the time T .

Proof:

We will begin with the case where $N \geq 3$ and we treat after the specific case $N = 2$.

Case $N \geq 3$:

We apply to the momentum equation the operator $(-\Delta)^{-1}\text{div}$ in order to concentrate us on the pressure and we get:

$$a\rho^\gamma = \frac{\partial}{\partial t}(-\Delta)^{-1}\text{div}(\rho u) + (-\Delta)^{-1}\partial_{i,j}^2(\rho u_i u_j) + (2\mu + \lambda)\text{div}u - \kappa(-\Delta)^{-1}\text{div}(\rho \nabla(\phi * \rho - \rho)), \quad (2.10)$$

and in multiplying by ρ^ε with $0 < \varepsilon \leq \min(\frac{1}{N}, \frac{2}{N}\gamma - 1)$ to estimate $\rho^{\gamma+\varepsilon}$, we get:

$$\begin{aligned} a\rho^{\gamma+\varepsilon} + \frac{\kappa}{2}\rho^{2+\varepsilon} &= -\kappa\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho(\nabla\phi * \rho)) + \rho^\varepsilon(-\Delta)^{-1}\partial_{i,j}^2(\rho u_i u_j) \\ &\quad + \frac{\partial}{\partial t}(\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)) - [\frac{\partial}{\partial t}\rho^\varepsilon](-\Delta)^{-1}\text{div}(\rho u) + (\mu + \zeta)\text{div}u, \end{aligned} \quad (2.11)$$

where we note $\xi = \lambda + \mu$. We now rewrite the previous equality as follows:

$$\begin{aligned} a\rho^{\gamma+\varepsilon} + \frac{\kappa}{2}\rho^{2+\varepsilon} &= -\kappa\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho(\nabla\phi * \rho)) + \rho^\varepsilon(-\Delta)^{-1}\partial_{i,j}^2(\rho(u_i)(u_j)) \\ &\quad + \frac{\partial}{\partial t}(\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)) + \text{div}[u\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)] + (\mu + \zeta)\text{div}u \\ &\quad - (\rho^\varepsilon u \cdot \nabla(-\Delta)^{-1}\text{div}(\rho u) + (1 - \varepsilon)(\text{div}u)\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)), \end{aligned} \quad (2.12)$$

Next we integrate (2.12) in time on $[0, T]$ and in space so we obtain:

$$\begin{aligned} \int_{(0,T) \times \mathbb{R}^N} (a\rho^{\gamma+\varepsilon} + \frac{\kappa}{2}\rho^{2+\varepsilon}) dx dt &= \int_{(0,T) \times \mathbb{R}^N} \left(\frac{\partial}{\partial t}[\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)] + (\mu + \zeta)(\text{div}u)\rho^\varepsilon \right. \\ &\quad + (1 - \varepsilon)(\text{div}u)\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u) + \rho^\varepsilon[R_i R_j(\rho u_i u_j) - u_i R_i R_j(\rho u_j)] \\ &\quad \left. + \text{div}[u\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)] - \kappa\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho \nabla(\phi * \rho)) \right) dx dt. \end{aligned} \quad (2.13)$$

where R_i is the classical Riesz transform.

Now we want to control the term $\int_0^T \int_{\mathbb{R}^N} (\rho^{\gamma+\varepsilon} + \frac{\kappa}{2}\rho^{2+\varepsilon}) dx dt$. As ρ is positive, it will enable us to control $\|\rho\|_{L_{t,x}^{\gamma+\varepsilon}}$ and $\|\rho\|_{L_{t,x}^{2+\varepsilon}}$. This may be achieved by controlling each term on the right side of (2.13).

We start by treating the term $\int_{(0,T) \times \mathbb{R}^N} \frac{\partial}{\partial t}[\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)]$. So we need to control $\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)$ in $L^\infty(0, T; L^1(\mathbb{R}^N))$ and $\rho_0^\varepsilon(-\Delta)^{-1}\text{div}(\rho_0 u_0)$ because:

$$\begin{aligned} \int_{(0,T) \times \mathbb{R}^N} \frac{\partial}{\partial t}[\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)](t, x) dt dx &= \int_{\mathbb{R}^N} \rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u)](x) dx(t) \\ &\quad - \int_{\mathbb{R}^N} \rho_0^\varepsilon(-\Delta)^{-1}\text{div}(\rho_0 u_0)](x) dx, \end{aligned}$$

We recall that ρ , ρ^2 , ρ^γ and $\rho|u|^2$ are bounded in $L^\infty(L^1)$ while Du is bounded in $L^2((0, T) \times \mathbb{R}^N)$ and u is bounded in $L^2(0, T; L^{\frac{2N}{N-2}}(\mathbb{R}^N))$ by Sobolev embedding. In

particular by Hölder inequalities we get that ρu is bounded in $L^\infty(0, T, (L^{\frac{2\gamma}{\gamma+1}} \cap L^{\frac{4}{3}})(\mathbb{R}^N))$.

Thus we get in using Hölder inequalities and Sobolev embedding:

$\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u) \in L^\infty(0, T, L^1 \cap L^\alpha)$ with:

$$\frac{1}{\alpha} = \frac{\varepsilon}{s} + \min\left(\frac{\gamma+1}{2\gamma}, \frac{3}{4}\right) - \frac{1}{N} < 1.$$

The fact that $\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho u) \in L^\infty(0, T, L^1)$ is obtained by interpolation because $\rho \in L^\infty(L^1)$ and in using less integrability in Sobolev embedding.

Next we have the same type of estimates for $\|\rho_0^\varepsilon(-\Delta)^{-1}\text{div}(\rho_0 u_0)\|_{L^1(\mathbb{R}^N)}$.

Finally (2.13) is rewritten on the following form in using Green formula:

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^N} (\rho^{\gamma+\varepsilon} + \frac{\kappa}{2} \rho^{2+\varepsilon}) dx dt &\leq C(1 + \int_0^T \int_{\mathbb{R}^N} [|\text{div} u| \rho^\varepsilon (1 + |(-\Delta)^{-1}\text{div}(\rho u)|) \\ &\quad + \rho^\varepsilon |R_i R_j(\rho u_i u_j) - u_i R_i R_j(\rho u_j)| + \kappa \rho^\varepsilon |(-\Delta)^{-1}\text{div}(\rho \nabla(\phi * \rho))|] dt dx). \end{aligned}$$

Now we will treat each term of the right hand side. We treat all the terms with the same type of estimates than P.-L. Lions in [15], excepted the capillarity term.

We start with the term $|\text{div} u| \rho^\varepsilon |(-\Delta)^{-1}\text{div}(\rho u)|$ where we have:

$$|\text{div} u| \in L^2(L^2), \quad \rho^\varepsilon \in L^\infty(L^{\frac{s}{\varepsilon}}), \quad \rho u \in L^2(0, T, L^r(\mathbb{R}^N))$$

with $\frac{1}{r} = \frac{1}{s} + \frac{N-2}{2N}$ and by Sobolev embedding $|(-\Delta)^{-1}\text{div}(\rho u)| \in L^2(L^{s'})$ with $\frac{1}{s'} = \frac{1}{r} - \frac{1}{N}$ (this is possible only if $r < N$). We are in a critical case for the Sobolev embedding (i.e $r \geq N$) only when $N = 3$ and $\gamma \geq 6$, that's why for $N = 3$ and $\gamma \geq 6$.

So by Hölder inequalities we get $|\text{div} u| \rho^\varepsilon |(-\Delta)^{-1}\text{div}(\rho u)| \in L^1(L^{s_1})$ with: $\frac{1}{s_1} = \frac{1}{s} + \frac{\varepsilon}{s} + \frac{1}{2} = 1 - \frac{2}{N} + \frac{1+\varepsilon}{s} \leq 1$ as we have $s > \frac{N}{2}$.

Moreover by interpolation $|\text{div} u| \rho^\varepsilon |(-\Delta)^{-1}\text{div}(\rho u)|$ belongs to $L^1(0, T; L^1(\mathbb{R}^N))$.

We now treat the case $N = 3$ and $\gamma \geq 6$ where we choose in this case $\varepsilon = \frac{2}{N}\gamma - 1$ to explicit precisely this case:

$$\begin{aligned} \| |\text{div} u| \rho^\varepsilon |(-\Delta)^{-1}\text{div}(\rho u)| \|_{L^1} &\leq \|Du\|_{L^2(L^2)} \|\rho\|_{L^{\gamma+\varepsilon}}^\varepsilon \|\rho u\|_{L^{\frac{2(\gamma+\varepsilon)}{\gamma-\varepsilon}}(L^{\frac{6(\gamma+\varepsilon)}{5\gamma-\varepsilon})}} \\ &\leq C \|\rho\|_{L^{\gamma+\varepsilon}}^\varepsilon \|\rho u\|_{L^{\frac{10\gamma-6}{\gamma+3}}(L^{\frac{3(10\gamma-6)}{13\gamma+3}})} \leq C \|\rho\|_{L^{\gamma+\varepsilon}}^\varepsilon \|\rho u\|_{L^2(L^{\frac{6\gamma}{\gamma+6}})}^{\frac{\gamma+3}{5\gamma-3}} \|\rho u\|_{L^\infty(L^2)}^{\frac{2(2\gamma-3)}{5\gamma-3}} \\ &\leq C \|\rho\|_{L^{\gamma+\varepsilon}}^\varepsilon \|\rho u\|_{L^2(L^{\frac{6\gamma}{\gamma+6}})}^{\frac{5\gamma}{5\gamma-3}} \|\rho u\|_{L^\infty(L^{\frac{2\gamma}{\gamma+1}})}^{\frac{2(2\gamma-3)}{5\gamma-3}} \\ &\leq C \|\rho\|_{L^{\gamma+\varepsilon}}^\varepsilon \end{aligned}$$

since we have $\frac{1}{2} + \frac{\varepsilon}{\gamma+\varepsilon} + \frac{\gamma-\varepsilon}{2(\gamma+\varepsilon)} = 1$, $\frac{1}{2} + \frac{\varepsilon}{\gamma+\varepsilon} + \frac{5\gamma-\varepsilon}{6(\gamma+\varepsilon)} - \frac{1}{3} = 1$, and $\frac{6(\gamma+\varepsilon)}{5\gamma-\varepsilon} = 3\frac{10\gamma-6}{13\gamma+3} < 3$.

We now want to treat the term: $\rho^\varepsilon |(-\Delta)^{-1}\text{div}(\rho \nabla(\phi * \rho))|$, so we have: $\rho \nabla(\phi * \rho) = \rho(\nabla \phi * \rho) \in L^\infty(L^1 \cap L^{\frac{s}{2}})$ by Hölder inequalities and the fact that we have $\rho \in L^\infty(L^1)$ and $\nabla \phi \in L^1$.

After we get that $\rho^\varepsilon(-\Delta)^{-1}\text{div}(\rho \nabla(\phi * \rho)) \in L^\infty(L^{r_1})$ with: $\frac{1}{r_1} = \frac{\varepsilon}{s} + \frac{2}{s} - \frac{1}{N} = \frac{2+\varepsilon}{s} - \frac{1}{N} < 1$.

We conclude that $\rho^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho\nabla(\phi*\rho))$ is $L^\infty(L^1)$ in using interpolation when $N = 2, 3$. Indeed we have $\rho\nabla(\phi*\rho) \in L^\infty(L^1)$ and in choosing $\varepsilon = \frac{2}{N}s - 1$ we have: $1 - \frac{1}{N} + \frac{2}{N}s - 1 \geq 1$ when $s \geq \frac{N}{2}(1 + \frac{1}{N})$. This is the case when $N = 2, 3$, and this is the case when $N \geq 4$ and $\gamma \geq \frac{N}{2}(1 + \frac{1}{N})$.

In the other case we need to work in arbitrary compact.

We have after the term $(\operatorname{div}(u))\rho^\varepsilon$. We recall that ρ^ε is in $L^\infty(L^{\frac{1}{\varepsilon}} \cap L^{\frac{s}{\varepsilon}})$. If $\varepsilon \geq \frac{1}{2}$ (i.e. $s \geq \frac{3}{4}N$), the bound is obvious because $\frac{1}{2} + \varepsilon \geq 1$ and $\frac{1}{2} + \frac{\varepsilon}{s} < 1$, we can then conclude by interpolation. On the other hand, this rather simple term presents a technical difficulty when $\varepsilon \leq \frac{1}{2}$ since we do not know in that case if $\operatorname{div}u\rho^\varepsilon \in L^1(\mathbb{R}^N \times (0, T))$. One way to get round the difficulty is to multiply (2.10) by $\rho^\varepsilon 1_{\{\rho \geq 1\}}$. Then we obtain an estimate on $\rho^{s+\varepsilon} 1_{\{\rho \geq 1\}}$ in $L^1((0, T) \times \mathbb{R}^N)$ as $\rho^\varepsilon 1_{\{\rho \geq 1\}} |\operatorname{div}u| \leq \rho |\operatorname{div}u| \in L^1((0, T) \times \mathbb{R}^N)$ (where $\varepsilon \leq \frac{1}{2}$) and we can conclude since $0 \leq \rho^{s+\varepsilon} 1_{\{\rho < 1\}} \leq \rho$ on $(0, T) \times \mathbb{R}^N$ and $\rho \in L^\infty(L^1)$.

We end with the following term $\rho^\varepsilon(R_i R_j(\rho u_i u_j) - u_i R_i R_j(\rho u_j))$. In the same way than in the previous inequalities we have $\rho^\varepsilon R_i R_j(\rho u_i u_j)$ is bounded in $L^1(0, T; L^1(\mathbb{R}^N))$. Indeed we have by Hölder inequalities and the fact that R_i is continuous from L^p in L^p with $1 < p < +\infty$: $\frac{1}{s} + 2\frac{N-2}{2N} + \frac{\varepsilon}{s} = 1 - \frac{2}{N} + \frac{1+\varepsilon}{s} \leq 1$ (because $s > \frac{N}{2}$). And we conclude by interpolation. We treat the term $\rho^\varepsilon u_i R_i R_j(\rho u_j)$ similarly.

We have to treat now the case $N = 2$ where we have to modify the estimates when we are in critical cases for Sobolev embedding.

Case $N = 2$:

In the case $N = 2$ most of the proof given above stay exact except for the slightly more delicate terms $\rho^\varepsilon |\operatorname{div}u|(-\Delta)^{-1}\operatorname{div}(\rho u)$ and $\rho^\varepsilon(R_i R_j(\rho u_i u_j) - u_i R_i R_j(\rho u_j))$.

We start with the term $|\rho^\varepsilon \operatorname{div}u(-\Delta)^{-1}\operatorname{div}(\rho u)|$. In our previous estimate it was possible to use Sobolev embedding on the term $(-\Delta)^{-1}\operatorname{div}(\rho u)$ only if $r \geq N$ (see above the notation), so in the case where $N = 2$ we are in a critical case for the Sobolev embedding when $\gamma \geq 2$.

This may be overcome by using that, by virtue of Sobolev embedding, we have:

$$\| |\operatorname{div}u| \rho^\varepsilon |(-\Delta)^{-1}\operatorname{div}(\rho u)| \|_{L^1} \leq C \|\rho\|_{L^{\gamma+\varepsilon}(L^{\gamma+\varepsilon})}^\varepsilon \|\rho u\|_{L^{2(\gamma+\varepsilon)}(L^{\frac{2(\gamma+\varepsilon)}{\gamma+\varepsilon+1}})}$$

Indeed: $\frac{1}{2} + \frac{\varepsilon}{\gamma+\varepsilon} + \frac{\gamma+\varepsilon+1}{2(\gamma+\varepsilon)} - \frac{1}{2} = \frac{1}{2} + \frac{2\varepsilon+1}{2\varepsilon+2\gamma} \leq 1 = \frac{1}{2} + \frac{\varepsilon}{\gamma+\varepsilon} + \frac{1}{2(\gamma+\varepsilon)} \leq 1$ and $\frac{1}{2} + \frac{\varepsilon}{\gamma+\varepsilon} + \frac{1}{2(\gamma+\varepsilon)} \leq 1$. Moreover we have as $\rho u = \sqrt{\rho}\sqrt{\rho}u$

$$\|\rho u\|_{L^{2(\gamma+\varepsilon)}(L^{\frac{2(\gamma+\varepsilon)}{\gamma+\varepsilon+1}})} \leq C \|\rho\|_{L^{\gamma+\varepsilon}(L^{\gamma+\varepsilon})}^{\frac{1}{2}}$$

then:

$$\| |\operatorname{div}u| \rho^\varepsilon |(-\Delta)^{-1}\operatorname{div}(\rho u)| \|_{L^1(L^1)} \leq C \|\rho\|_{L^{\gamma+\varepsilon}(L^{\gamma+\varepsilon})}^{\varepsilon+\frac{1}{2}}.$$

Next we are interested by the term $\rho^\varepsilon(R_i R_j(\rho u_i u_j) - u_i R_i R_j(\rho u_j))$. We use the fact that u is bounded in $L^2(0, T; \dot{H}^1)$ and thus in $L^2(0, T; BMO)$. Then by the Coifman-Rochberg-Weiss commutator theorem in [3], we have for almost all $t \in [0, T]$:

$$\|R_i R_j(\rho u_i u_j) - u_i R_i R_j(\rho u_j)\|_{L^{\frac{2(\gamma+\varepsilon)}{\gamma+\varepsilon+1}}(L^{\frac{2(\gamma+\varepsilon)}{\gamma+\varepsilon+1}})} \leq C \|u\|_{L^2(BMO)} \|\rho u\|_{L^{2(\gamma+\varepsilon)}(L^{\frac{2(\gamma+\varepsilon)}{\gamma+\varepsilon+1}})}$$

So we have:

$$\|\rho^\varepsilon(R_i R_j(\rho u_i u_j) - u_i R_i R_j(\rho u_j))\|_{L^1} \leq C \|\rho\|_{L^{\gamma+\varepsilon}(L^{\gamma+\varepsilon})}^{\varepsilon+\frac{1}{2}}$$

In view of the previous inequalities we get finally:

$$\|\rho\|_{L^{\gamma+\varepsilon}(L^{\gamma+\varepsilon})}^{\gamma+\varepsilon} \leq C(1 + \|\rho\|_{L^{\gamma+\varepsilon}(L^{\gamma+\varepsilon})}^{\frac{1}{2}+\varepsilon})$$

and the $L^{\gamma+\varepsilon}(L^{\gamma+\varepsilon})$ bound on ρ is proven since $\frac{1}{2} + \varepsilon < \gamma + \varepsilon$. $\square$

2.2 Compactness results for compressible Navier-Stokes equations of Korteweg type in the case of isentropic pressure

In the sequel we are not going to treat in details the case with $N \geq 4$ and $\gamma < \frac{N}{2}(1 + \frac{1}{N})$, we just remark that the proof is the same as in the case $N = 2$, it suffices to localize because we can only apply the theorem 2.2 on the gain of integrability on any compact K .

So let follow the theorem 2.2 and assume that $\gamma > \frac{N}{2}$ if $N \geq 4$ and $\gamma \geq 1$ such that if (ρ, u) is a regular solution then $\rho \in L^q((0, T) \times \mathbb{R}^N)$ with $q = \gamma + 1 - \frac{2\gamma}{N}$. We can observe that in this case $q > s = \max(\gamma, 2)$. We will see that it will be very useful in the sequel to justify the passage to the weak limit in some terms to get a gain of integrability on the density. Indeed the key point to proving the existence of weak solutions is the passage to the limit in the term of pressure and in the term of capillarity $\rho \nabla(\phi * \rho - \rho)$.

First, we assume that a sequence $(\rho_n, u_n)_{n \in \mathbb{N}}$ of approximate weak solutions has been constructed by a mollifying process, which have suitable regularity to justify the formal estimates like the energy estimate and the theorem 2.2. $(\rho_n, u_n)_{n \in \mathbb{N}}$ has the initial data of the theorem 1.1 with uniform bounds, i.e:

Moreover $(\rho_n, u_n)_{n \in \mathbb{N}}$ verifies the energy inequalities (1.4) and the previous theorem 2.2, we have then:

- ρ_n is bounded uniformly in $L^\infty(0, T; L^1 \cap L^s(\mathbb{R}^N)) \cap C([0, T]; L^p(\mathbb{R}^N))$ for $1 \leq p < \max(2, \gamma)$,
- $\rho_n \geq 0$ a.e. ρ_n is bounded uniformly in $L^q(0, T; \mathbb{R}^N)$ with $q > s$,
- ∇u_n is bounded in $L^2(0, T; L^2(\mathbb{R}^N))$, $\rho_n |u_n|^2$ is bounded in $L^\infty(0, T; L^1(\mathbb{R}^N))$,
- u_n is bounded in $L^2(0, T; L^{\frac{2N}{N-2}}(\mathbb{R}^N))$ for $N \geq 3$.

And we have in passing to the weak limit in the previous bound in extracting subsequence if necessary:

- $\rho_n \rightharpoonup \rho$ weakly in $L^s((0, T) \times \mathbb{R}^N)$,

- $u_n \rightarrow u$ weakly in $L^2(0, T, \dot{H}^1(\mathbb{R}^N))$,
- $\rho_n^\gamma \rightarrow \overline{\rho^\gamma}$ weakly in $L^r((0, T) \times \mathbb{R}^N)$ for $r = \frac{q}{\gamma} > 1$,
- $\rho_n^2 \rightarrow \overline{\rho^2}$ weakly in $L^{r_1}((0, T) \times \mathbb{R}^N)$ for $r_1 = \frac{q}{2} > 1$.

Notation 1 We will always write in the sequel $\overline{B(\rho)}$ to mean the weak limit of the sequence $B(\rho_n)$ bounded in appropriate space that we will precise.

We recall that the main difficulty will be to pass to the limit in the pressure term and the capillary term. The idea of the proof will be to test the convergence of the sequence $(\rho_n)_{n \in \mathbb{N}}$ on convex functions B in order to use their properties of lower semi-continuity with respect to the weak topology in $L^1(\mathbb{R}^N)$. In this goal we will use the theory of renormalized solutions introduced by Diperna and Lions in [?]. So we will obtain strong convergence of ρ_n in appropriate spaces.

2.3 Idea of the proof

We here give a sketchy proof of the theorem 1.1. In this spirit we can rewrite mass conservation of the regular solution $(\rho_n, u_n)_{n \in \mathbb{N}}$ on the form:

$$\frac{p}{\partial t}(B(\rho_n)) + \operatorname{div}(u_n B(\rho_n)) = (B(\rho_n) - \rho_n B'(\rho_n)) \operatorname{div} u_n.$$

Supposing that $B(\rho_n)$ is bounded in appropriate space we can pass to the weak limit where we have in the energy space $\rho_n \rightharpoonup \rho$ and $u_n \rightharpoonup u$, so we get:

$$\frac{p}{\partial t}(\overline{B(\rho)}) + \operatorname{div}(u \overline{B(\rho)}) = \overline{(B(\rho) - \rho B'(\rho)) \operatorname{div} u} \quad (2.14)$$

We will set: $b(\rho) = B(\rho) - \rho B'(\rho)$.

Next in seeing the mass equation for approximate solutions $(\rho_n, u_n)_{n \in \mathbb{N}}$, and in passing directly to the limit via the weak convergence arguing like P-L. Lions in [15] p 13 we get:

$$\frac{d}{dt} \rho + \operatorname{div}(\rho u) = 0. \quad (2.15)$$

After we will just have to verify the passage to the limit for the product ρu . Next we will use the theorem on the renormalized solutions of Diperna-Lions in [14] on (2.15) in recalling that $\rho \in L^\infty(L^2)$. So we get:

$$\frac{d}{dt}(B(\rho)) + \operatorname{div}(u B(\rho)) = b(\rho) \operatorname{div}(u). \quad (2.16)$$

Next we subtract (2.14) to (2.16), so we obtain:

$$\frac{d}{dt}(\overline{B(\rho)} - B(\rho)) + \operatorname{div}(u(\overline{B(\rho)} - B(\rho))) = \overline{b(\rho) \operatorname{div} u} - b(\rho) \operatorname{div} u. \quad (2.17)$$

Consequently, in order to estimate the difference $\overline{B(\rho)} - B(\rho)$ which tests the convergence of ρ_n , we need to estimate the difference $\overline{b(\rho) \operatorname{div}(u)} - b(\rho) \operatorname{div}(u)$. We choose then B a concave function and we have:

$$\overline{B(\rho)} - B(\rho) \leq 0.$$

The goal will be now to prove the reverse inequality in order to justify that $B(\rho_n)$ tends to $B(\rho)$ a.e.

So now we aim at estimating the difference $\overline{b(\rho)\operatorname{div}u} - b(\rho)\operatorname{div}u$. This may be achieved by introducing the effective viscous pressure $P_{eff} = P - (2\mu + \lambda)\operatorname{div}u$ after D. Hoff in [?], which satisfies some important properties of weak convergence.

In fact owing to the capillarity term we adapt Hoff's concept to our equation in setting:

$$\tilde{P}_{eff} = P + \frac{\kappa}{2}\rho^2 - (2\mu + \lambda)\operatorname{div}u.$$

Proof of theorem 1.1

We begin with the case $N \geq 3$, and next we will complete the proof by the case $N = 2$ in specifying the changes to bring.

Before getting into the heart of the proof, we first recall that we obtain easily the convergence in distribution sense of $\rho_n u_n$ to ρu and $\rho_n(u_n)_i(u_n)_j$ to $\rho u_i u_j$. We refer to the classical result by Lions (see [15]) or the book of Novotný and Straškraba [16].

Case $N \geq 3$

We have seen in the idea of the proof 2.3 that our goal is to compare $\overline{B(\rho)}$ and $B(\rho)$ for certain concave functions B . From the mass equation we have obtained:

$$\partial_t(\overline{B(\rho)} - B(\rho)) + \operatorname{div}(u(\overline{B(\rho)} - B(\rho))) = \overline{b(\rho)\operatorname{div}(u)} - b(\rho)\operatorname{div}(u). \quad (2.18)$$

So before comparing $\overline{B(\rho)}$ and $B(\rho)$, we have to investigate the expression $\overline{b(\rho)\operatorname{div}(u)} - b(\rho)\operatorname{div}(u)$. By virtue of theorem 2.2 which gives a gain of integrability we can take the function $B(x) = x^\varepsilon$, as we control for ε small enough $\rho^{s+\varepsilon}$. Our goal now is to exhibit the effective pressure $\tilde{P}_{eff}$, and to multiply it by ρ^ε to extract $\overline{\operatorname{div}u b(\rho)}$. We will see in the sequel how to compare it with $b(\rho)\operatorname{div}(u)$. So we focus on the convergence of the pressure and capillarity terms.

Control of the term $\overline{\operatorname{div}u b(\rho)}$

So we take the div on the momentum equation satisfied by the regular solution. We get:

$$\frac{\partial}{\partial t}\operatorname{div}(\rho_n u_n) + \partial_{ij}^2(\rho_n u_n^i u_n^j) - \zeta \Delta \operatorname{div}u_n + \Delta(a\rho_n^\gamma) = \kappa \operatorname{div}(\rho_n(\nabla\phi * \rho_n)) - \frac{\kappa}{2}\Delta(\rho_n^2), \quad (2.19)$$

with $\zeta = \lambda + 2\mu$. Applying the operator $(-\Delta)^{-1}$ to (2.19), we obtain:

$$\begin{aligned} \frac{\partial}{\partial t}(-\Delta)^{-1}\operatorname{div}(\rho_n u_n) + (-\Delta)^{-1}\partial_{ij}^2(\rho_n u_n^i u_n^j) + [\zeta \operatorname{div}u_n - a\rho_n^\gamma - \frac{\kappa}{2}\rho_n^2] \\ = \kappa(-\Delta)^{-1}\operatorname{div}(\rho_n(\nabla\phi * \rho_n)). \end{aligned} \quad (2.20)$$

After we multiply (2.20) by ρ_n^ε with ε that we choose enough small with $\varepsilon \in (0, 1)$:

$$\begin{aligned} [(\mu + \zeta)\operatorname{div}u_n - a\rho_n^\gamma - \frac{\kappa}{2}\rho_n^2]\rho_n^\varepsilon = \kappa\rho_n^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n(\nabla\phi * \rho_n)) \\ - \rho_n^\varepsilon \frac{\partial}{\partial t}(-\Delta)^{-1}\operatorname{div}(\rho_n u_n) - \rho_n^\varepsilon(-\Delta)^{-1}\partial_{ij}^2(\rho_n u_n^i u_n^j). \end{aligned} \quad (2.21)$$

So if we rewrite (2.21), we have:

$$[(\mu + \zeta)\operatorname{div} u_n - a\rho_n^\gamma - \frac{\kappa}{2}\rho_n^2]\rho_n^\varepsilon = \kappa\rho_n^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n(\nabla\phi * \rho_n)) - \rho_n^\varepsilon(-\Delta)^{-1}\partial_{ij}^2(\rho_n u_n^i u_n^j) \\ - \frac{\partial}{\partial t}((\rho_n)^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n u_n)) + [\frac{\partial}{\partial t}(\rho_n)^\varepsilon](-\Delta)^{-1}\operatorname{div}(\rho_n u_n),$$

Next we have:

$$[(\mu + \zeta)\operatorname{div} u_n - a\rho_n^\gamma - \frac{\kappa}{2}\rho_n^2]\rho_n^\varepsilon = \kappa\rho_n^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n(\nabla\phi * \rho_n)) - \rho_n^\varepsilon(-\Delta)^{-1}\partial_{ij}^2(\rho_n u_n^i u_n^j) \\ - \frac{\partial}{\partial t}[(\rho_n)^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n u_n)] - \operatorname{div}[u_n(\rho_n)^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n u_n)] \\ + (\rho_n)^\varepsilon u_n \cdot \nabla(-\Delta)^{-1}\operatorname{div}(\rho_n u_n) + (1 - \varepsilon)(\operatorname{div} u_n)(\rho_n)^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n u_n), \quad (2.22)$$

or finally:

$$[(\mu + \xi)\operatorname{div} u_n - a\rho_n^\gamma - \frac{\kappa}{2}\rho_n^2]\rho_n^\varepsilon = \kappa\rho_n^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n(\nabla\phi * \rho_n)) \\ - \frac{\partial}{\partial t}[\rho_n^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n u_n)] - \operatorname{div}[u_n(\rho_n)^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n u_n)] \quad (2.23) \\ + (\rho_n)^\varepsilon[u_n \cdot \nabla(-\Delta)^{-1}\operatorname{div}(\rho_n u_n) - (-\Delta)^{-1}\partial_{ij}^2(\rho u_n^i u_n^j)] \\ + (1 - \varepsilon)(\operatorname{div} u_n)(\rho_n)^\varepsilon(-\Delta)^{-1}\operatorname{div}(\rho_n u_n) .$$

Now like in Lions [?] we want to pass to the limit in the distribution sense in (2.23) in order to estimate $\overline{\operatorname{div} u(\rho)^\varepsilon}$.

Passage to the weak limit in (2.23)

In this goal we use the following lemma of P-L Lions in [15] to express the weak limit of the non-linear terms.

Lemma 1 *Let Ω a open of $\mathbb{R}^N$. Let (g_n, h_n) converge weakly to (g, h) in $L^{p_1}(0, T, L^{p_2}(\Omega)) \times L^{q_1}(0, T, L^{q_2}(\Omega))$ where $1 \leq p_1, p_2, q_1, q_2 \leq +\infty$ satisfy,*

$$\frac{1}{p_1} + \frac{1}{q_1} = \frac{1}{p_2} + \frac{1}{q_2} = 1 .$$

We assume in addition that:

$$\frac{\partial g^n}{\partial t} \text{ is bounded in } L^1(0, T, W^{-m, 1}(\Omega)) \text{ for some } m \geq 0 \text{ independent of } n. \quad (2.24)$$

and that:

$$\|h^n - h^n(\cdot, \cdot + \xi)\|_{L^{q_1}(0, T, L^{q_2}(\Omega))} \rightarrow 0 \quad \text{as } |\xi| \rightarrow 0, \text{ uniformly in } n. \quad (2.25)$$

Then, $g^n h^n$ converges to gh (in the sense of distribution on $\Omega \times (0, T)$).

So we use the above lemma to pass to the weak limit in the four following non-linear terms of (2.23):

$$\begin{aligned} T_n^1 &= u_n \rho_n^\varepsilon (-\Delta)^{-1} \operatorname{div}(\rho_n u_n), & T_n^2 &= \rho_n^\varepsilon (-\Delta)^{-1} \operatorname{div}(\rho_n u_n), \\ T_n^3 &= \rho_n^\varepsilon (-\Delta)^{-1} \operatorname{div}(\rho_n (\nabla \phi * \rho_n)), & T_n^4 &= (\operatorname{div} u_n) (\rho_n)^\varepsilon (-\Delta)^{-1} \operatorname{div}(\rho_n u_n). \end{aligned}$$

So we choose the different g_n^i and h_n^i as follows:

$$\begin{aligned} \text{for } T_n^1 & \quad g_n^1 = u_n (\rho_n)^\varepsilon & g^1 &= u \overline{\rho^\varepsilon} & h_n^1 &= (-\Delta)^{-1} \operatorname{div}(\rho_n u_n) \\ \text{for } T_n^2 & \quad g_n^2 = \rho_n^\varepsilon & g^2 &= \overline{\rho^\varepsilon} & h_n^2 &= (-\Delta)^{-1} \operatorname{div}(\rho_n u_n) \\ \text{for } T_n^3 & \quad g_n^3 = \rho_n^\varepsilon & g^3 &= \overline{\rho^\varepsilon} & h_n^3 &= (-\Delta)^{-1} \operatorname{div}(\rho_n (\nabla \phi * \rho_n)) \\ \text{for } T_n^4 & \quad g_n^4 = (\operatorname{div} u_n) (\rho_n)^\varepsilon & g^4 &= \overline{\operatorname{div} u} \overline{\rho^\varepsilon} & h_n^4 &= (-\Delta)^{-1} \operatorname{div}(\rho_n u_n). \end{aligned}$$

To show that $u_n (\rho_n)^\varepsilon$ converges in distribution sense to $u \overline{\rho^\varepsilon}$ we apply easily lemma 1 with $h_n = u_n$ and $g_n = \rho_n^\varepsilon$. We now want to examine each term and apply the above lemma to pass to the limit in the weak sense.

We start with the first term T_n^1 . We have that $\rho_n^\varepsilon u_n \in L^\infty(L^q) \cap L^2(L^r)$ with $\frac{1}{q} = \frac{\varepsilon}{2s} + \frac{1}{2}$ and $\frac{1}{r} = \frac{(N-2)}{2N} + \frac{\varepsilon}{s} = \frac{1}{2} - \frac{1}{N} + \frac{\varepsilon}{s}$. In addition the hypothesis (2.24) is immediately verified (use the momentum equation).

We now want to verify the hypothesis (2.25), so we have h_n^1 belongs to $L^\infty(W_{loc}^{1,q'}(\mathbb{R}^N)) \cap L^2(W_{loc}^{1,r'}(\mathbb{R}^N))$ with $\frac{1}{q'} = \frac{1}{2} + \frac{1}{2s}$ and $\frac{1}{r'} = \frac{(N-2)}{2N} + \frac{1}{s} = \frac{1}{2} + \frac{1}{s} - \frac{1}{N}$. This result enables us to verify the hypothesis (2.25) by Sobolev embedding.

So we can choose (with the notation of the above lemma) $q_1 = 2$ and $q_2 \in (r', \frac{Nr'}{N-r'})$, $p_1 = 2$, and $p_2 = 1 - \frac{1}{q_2}$ which is possible by interpolation. Indeed we have: $\frac{1}{r'} + \frac{1}{r} = 1 - \frac{2}{N} + \frac{1+\varepsilon}{s} \leq 1$.

We proceed in the same way for T_n^2 and T_n^4 .

We can similarly examine T_n^3 , because $\rho_n^\varepsilon \in L^\infty(L^{\frac{1}{\varepsilon}} \cap L^{\frac{s}{\varepsilon}})$ and $\rho_n (\nabla \phi * \rho_n) \in L^\infty(L^1 \cap L^{\frac{s}{2}})$, we can choose $p_2 = \frac{1}{\varepsilon}$, we have then $(\Delta)^{-1} \operatorname{div} \rho_n (\nabla \phi * \rho_n) \in L^\infty(0, T; W^{1, \frac{s}{2}})$ so that we can choose $q_1 = 2$, $q_2 \in (1, \frac{N \frac{s}{2}}{N - \frac{s}{2}})$. We can conclude by interpolation.

Finally we have to study the last non linear following term that we treat similarly as P-L.Lions in [15]:

$$A_n = (\rho_n)^\varepsilon [u_n \cdot \nabla (-\Delta)^{-1} \operatorname{div}(\rho_n u_n) - (-\Delta)^{-1} \partial_{ij}^2 (\rho_n (u_i)_n (u_j)_n)].$$

We can express this term A_n as follows:

$$A_n = (\rho_n)^\varepsilon [u_n^j, R_{ij}] (\rho_n u_n^i).$$

where $R_{ij} = (-\Delta)^{-1} \partial_{ij}^2$ with R_i the classical Riesz transform.

Next, we use a result by Coifman, Meyer on this type of commutator (see [5]) to take advantage of the regularity of $[u_n^j, R_{ij}] (\rho_n u_n^i)$.

Theorem 2.3 *The following map is continuous for any $N \geq 2$:*

$$\begin{aligned} W^{1,r_1}(\mathbb{R}^N)^N \times L^{r_2}(\mathbb{R}^N) &\rightarrow W^{1,r_3}(\mathbb{R}^N)^N \\ (a, b) &\rightarrow [a_j, R_i R_j] b_i \end{aligned} \tag{2.26}$$

with: $\frac{1}{r_3} = \frac{1}{r_1} + \frac{1}{r_2}$.

To pass to the weak limit in A_n we will use the previous lemma. We start with the case with $s > 3$. This quantity belongs to the space $L^1(W^{1,q})$ provided that $Du_n \in L^2(L^2)$ and $\rho u^j \in L^2(L^r)$ where $\frac{1}{r} = \frac{N-2}{2N} + \frac{1}{s} = \frac{1}{2} - \frac{1}{N} + \frac{1}{s}$ in which case $\frac{1}{q} = \frac{1}{r} + \frac{1}{2} = 1 - \frac{1}{N} + \frac{1}{s} \leq 1$. After we can use the above lemma applied to $h_n = [R_{ij}, u_n^j](\rho_n u_n^i)$ and $g_n = \rho_n^\varepsilon$. We can show easily in using again lemma 1 that h_n converges in distribution sense to $[R_{ij}, u_j](\rho_n u_i)$.

So we can take: $q_1 = 1$, $p_1 = +\infty$ and $q_2 \in (q, \frac{qN}{N-q})$, $p_2 = 1 - \frac{1}{q_2}$, this one because we can use interpolation and we can localize as we want limit in distribution sense.

In the case where $s \leq 3$, a simple interpolation argument can be used to accommodate the general case. It suffices to fix $L^{r_2}(\mathbb{R}^N)$ in the application (2.26) and use a result of Riesz-Thorin.

Finally according to the equation (2.21), and after passing to the limit we get:

$$\begin{aligned} [(\mu + \xi) \overline{\text{div} u \rho^\varepsilon} - \overline{(a\rho^{\gamma+\varepsilon})} - \frac{\kappa}{2} \overline{\rho^{2+\varepsilon}}] &= \overline{\rho^\varepsilon}(-\Delta)^{-1} \text{div}(\overline{\rho(\nabla \phi * \rho)}) - \frac{\partial}{\partial t} [\overline{\rho^\varepsilon}(-\Delta)^{-1} \text{div}(\rho u)] \\ &\quad - \text{div}[\overline{\rho^\varepsilon} u (-\Delta)^{-1} \text{div}(\rho u)] + \overline{\rho^\varepsilon} [u \cdot \nabla (-\Delta)^{-1} \text{div}(\rho u) - (-\Delta)^{-1} \partial_{ij}(\rho u_i u_j)] \\ &\quad + (1 - \varepsilon) \overline{\text{div} u \rho^\varepsilon} (-\Delta)^{-1} \text{div}(\rho u). \end{aligned} \quad (2.27)$$

Inequality between the terms $\overline{\rho^\varepsilon} \text{div} u$ and $\overline{\text{div} u \rho^\varepsilon}$

Now we are interested in estimating the term $\overline{\rho^\varepsilon} \text{div} u$ in order to describe the quantity $\overline{\rho^\varepsilon} \text{div} u - \overline{\text{div} u \rho^\varepsilon}$ before considering the quantity $\rho^\varepsilon \text{div} u - \overline{\text{div} u \rho^\varepsilon}$. We pass to the weak limit directly in (2.20) and we get in using again the lemma ??:

$$\begin{aligned} \frac{\partial}{\partial t} (-\Delta)^{-1} \text{div}(\rho u) + (-\Delta)^{-1} \partial_{ij}^2(\rho u_i u_j) + [(\mu + \xi) \text{div} u - \overline{a\rho^\gamma}] &= \\ &= -(-\Delta)^{-1} \text{div}(\overline{\rho(\nabla \phi * \rho)}) + \frac{\kappa}{2} \overline{\rho^2}. \end{aligned} \quad (2.28)$$

Now we just multiply (2.28) with $\overline{\rho^\varepsilon}$ and we can see that each term has a distribution sense.

So we get in proceeding in the same way as before:

$$\begin{aligned} [(\mu + \xi) \text{div} u \overline{\rho^\varepsilon} - \overline{(a\rho^\gamma)} \overline{\rho^\varepsilon} - \frac{\kappa}{2} \overline{\rho^2 \rho^\varepsilon}] &= \overline{\rho^\varepsilon}(-\Delta)^{-1} \text{div}(\overline{\rho(\nabla \phi * \rho)}) \\ &\quad - \overline{\rho^\varepsilon} \frac{\partial}{\partial t} [\rho (-\Delta)^{-1} \text{div}(\rho u)] + \overline{\rho^\varepsilon} [u \cdot \nabla (-\Delta)^{-1} \text{div}(\rho u) - (-\Delta)^{-1} \partial_{ij}(\rho u_i u_j)] \\ &\quad - \text{div}[\rho^\varepsilon u (-\Delta)^{-1} \text{div}(\rho u)] + (1 - \varepsilon) \overline{\text{div} u (\rho)^\varepsilon} (-\Delta)^{-1} \text{div}(\rho u). \end{aligned} \quad (2.29)$$

In subtracting (2.29) from (2.27), we get:

$$(\mu + \xi) \overline{\text{div} u (\rho)^\varepsilon} - \overline{a\rho^{\gamma+\varepsilon}} - \frac{\kappa}{2} \overline{\rho^{2+\varepsilon}} = (\mu + \xi) \text{div} u \overline{\rho^\varepsilon} - \overline{a\rho^\gamma} \overline{\rho^\varepsilon} - \frac{\kappa}{2} \overline{\rho^2 \rho^\varepsilon} \quad \text{a.e.}$$

Next we observe that by convexity:

$$(\overline{\rho^{\gamma+\varepsilon}})^{\frac{\varepsilon}{\gamma+\varepsilon}} \geq (\overline{\rho^\varepsilon}), \quad (\overline{\rho^{\gamma+\varepsilon}})^{\frac{\gamma}{\gamma+\varepsilon}} \geq (\overline{\rho^\gamma}) \quad \text{a.e.}$$

So we get:

$$\overline{\operatorname{div} u(\rho)^\varepsilon} \geq \operatorname{div} u \overline{\rho^\varepsilon}. \quad (2.30)$$

Comparison between ρ and $\overline{\rho^\varepsilon}^{\frac{1}{\varepsilon}}$

As on since (ρ_n, u_n) are regular solutions we get the equality (2.16) applied to $B(x) = x^\varepsilon$. So we get:

$$\frac{\partial}{\partial t}(\rho_n)^\varepsilon + \operatorname{div}(u_n(\rho_n)^\varepsilon) = (1 - \varepsilon)\operatorname{div} u_n(\rho_n)^\varepsilon. \quad (2.31)$$

And after we pass to the weak limit in (2.31) and we get:

$$\frac{\partial}{\partial t} \overline{\rho^\varepsilon} + \operatorname{div}(u \overline{\rho^\varepsilon}) = (1 - \varepsilon)\overline{\operatorname{div} u \rho^\varepsilon}. \quad (2.32)$$

When we pass to the weak limit in combining with (2.30) we get:

$$\frac{\partial}{\partial t} \overline{(\rho)^\varepsilon} + \operatorname{div}(u \overline{(\rho)^\varepsilon}) \geq (1 - \varepsilon)\overline{\operatorname{div} u (\rho)^\varepsilon}. \quad (2.33)$$

Now we wish to conclude about the pointwise convergence of ρ_n in proving that $(\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}} = \rho$ and to finish we will use the following theorem (see [8] p 34) applied to $B(x) = x^{\frac{1}{\varepsilon}}$ which is convex.

Theorem 2.4 *Let $(v_n)_{n \in \mathbb{N}}$ be a sequence of functions bounded in $L^1(\mathbb{R}^N)$ such that:*

$$v_n \rightharpoonup v \quad \text{weakly in } L^1(\mathbb{R}^N).$$

Let $\varphi : \mathbb{R} \rightarrow (-\infty, +\infty]$ be a upper semi-continuous strictly concave function such that $\varphi(v_n) \in L^1(\mathbb{R}^N)$ for any n , and:

$$\varphi(v_n) \rightharpoonup \overline{\varphi(v)} \quad \text{weakly in } L^1(\mathbb{R}^N).$$

Then:

$$v_n(y) \rightarrow v(y) \quad \text{a.e.}$$

extracting a subsequence as the case may be.

Now we want to use a type of Diperna-Lions theorem on inequality (2.33). Our goal is to renormalize this inequality with the function $B(x) = x^{\frac{1}{\varepsilon}}$ so that one can compare ρ and $\overline{\rho^\varepsilon}^{\frac{1}{\varepsilon}}$. Although (2.33) doesn't correspond exactly to the mass equation, we can use the same technics to renormalize the solution provided that $\rho \in L^\infty(L^2)$ which is the case. In our case it is very important that $\rho \in L^\infty(L^2)$, indeed it avoids to have supplementary conditions on the index γ like for the compressible Navier-Stokes system in [15]. We recall of Diperna-Lions theorem on renormalized solution for the mass equation.

Theorem 2.5 *Suppose that $\rho \in L^\infty(L^2)$ and $\beta \in C^1([0, \infty); \mathbb{R})$.*

We have then:

$$\frac{\partial \beta(\rho)}{\partial t} + \operatorname{div}(\beta(\rho) u) = (\beta(\rho) - \rho \beta'(\rho)) \operatorname{div} u$$

in distribution sense.

We now want to adapt this theorem for our equation (??) with $\beta(x) = x^{\frac{1}{\varepsilon}}$, so we may regularize by ω_α (with $\omega_\alpha = \frac{1}{\alpha^N} \omega(\frac{\cdot}{\alpha})$ where $\omega \in C_0^\infty(\mathbb{R}^N)$, $\text{supp } \omega \in B_1$ and $\int \omega dx = 1$) and find for all $\beta \in C_0^\infty([0, +\infty))$:

$$\frac{\partial}{\partial t}(\overline{\rho^\varepsilon} * \omega_\alpha) + \text{div}[u \overline{\rho^\varepsilon} * \omega_\alpha] \geq (1 - \varepsilon) \text{div} u \overline{\rho^\varepsilon} * \omega_\alpha + R_\alpha$$

where we have:

$$\begin{aligned} R_\alpha &= \text{div}[u \overline{\rho^\varepsilon} * \omega_\alpha] - \text{div}(u \overline{\rho^\varepsilon}) * \omega_\alpha + (1 - \varepsilon)[\text{div} u \overline{\rho^\varepsilon}] * \omega_\alpha - (1 - \varepsilon) \text{div} u \overline{\rho^\varepsilon} * \omega_\alpha \\ \frac{\partial}{\partial t}(\beta(\overline{\rho^\varepsilon} * \omega_\alpha)) + \text{div}[u \beta(\overline{\rho^\varepsilon} * \omega_\alpha)] &\geq (1 - \varepsilon) \text{div} u \overline{\rho^\varepsilon} * \omega_\alpha \beta'(\overline{\rho^\varepsilon} * \omega_\alpha) \\ &+ (\text{div} u)[\beta(\overline{\rho^\varepsilon} * \omega_\alpha) - \overline{\rho^\varepsilon} * \omega_\alpha \beta'(\overline{\rho^\varepsilon} * \omega_\alpha)] + R_\alpha \beta'(\overline{\rho^\varepsilon} * \omega_\alpha) \\ &= -\varepsilon(\text{div} u) \overline{(\rho)^\varepsilon} \beta'(\overline{\rho^\varepsilon}) + (\text{div} u) \beta(\overline{\rho^\varepsilon}) + R_\alpha \beta'(\overline{\rho^\varepsilon} * \omega_\alpha). \end{aligned}$$

After we pass to the limit when $\alpha \rightarrow 0$ and we see that R_α tends to 0 in using lemma on regularization in [14] p 43. This looks like a rather innocent manipulation but it's at this point that we require to control ρ in $L^2(0, T; \mathbb{R}^N)$. And in our case we don't need to impose $\gamma > \frac{N}{2}$ for $N = 2, 3$. Hence:

$$\frac{\partial}{\partial t}(\beta(\overline{(\rho)^\varepsilon})) + \text{div}[u \beta(\overline{(\rho)^\varepsilon})] \geq -\varepsilon(\text{div} u) \overline{\rho^\varepsilon} \beta'(\overline{\rho^\varepsilon}) + (\text{div} u) \beta(\overline{\rho^\varepsilon}).$$

We then choose $\beta = (\Psi_M)^{\frac{1}{\varepsilon}}$ where $\Psi_M = M \Psi(\frac{\cdot}{M})$, $M \geq 1$, $\Psi \in C_0^\infty([0, +\infty))$, $\Psi(x) = x$ on $[0, 1]$, $\text{supp } \Psi \subset [0, 2]$, and we obtain:

$$\begin{aligned} \frac{\partial}{\partial t}(\Psi_M(\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}}) + \text{div}[u \Psi_M(\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}}] &\geq (\text{div} u) \Psi_M(\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}-1} \Psi'_M(\overline{\rho^\varepsilon}) \overline{\rho^\varepsilon} + (\text{div} u) \Psi_M(\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}} \\ &\geq \text{div} u \Psi_M(\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}-1} [\Psi_M(\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}} - \Psi'_M(\overline{\rho^\varepsilon}) \overline{\rho^\varepsilon}] 1_{(\overline{\rho^\varepsilon} > M)} \\ &\geq -C_0 |\text{div} u| M^{\frac{1}{\varepsilon}} 1_{(\overline{\rho^\varepsilon} > M)}. \end{aligned}$$

where $C_0 = \sup\{|\Psi(x)|^{\frac{1}{\varepsilon}-1} |\Psi(x) - x \Psi'(x)|, x \in [0, +\infty)\}$.

Now we claim that:

$$\frac{\partial}{\partial t}(\overline{(\rho)^\varepsilon})^{\frac{1}{\varepsilon}} + \text{div}(u \overline{(\rho)^\varepsilon})^{\frac{1}{\varepsilon}} \geq 0. \quad (2.34)$$

For proving that, we notice that by convexity $\overline{(\rho)^\varepsilon}^{\frac{1}{\varepsilon}} \leq \rho$, so we get :

$$\| |\text{div} u| M^{\frac{1}{\varepsilon}} 1_{(\overline{\rho^\varepsilon} > M)} \|_{L_T^1(L^1(\mathbb{R}^N))} \leq \| \text{div} u \|_{L_T^2(L^2(\mathbb{R}^N))} \| \rho 1_{\rho > M^{\frac{1}{\varepsilon}}} \|_{L_T^2(L^2(\mathbb{R}^N))} \rightarrow 0 \text{ as } M \rightarrow +\infty.$$

We have concluded by dominated convergence. We have concluded by dominated convergence.

At this stage we subtract the mass equation to (2.34) and we get in setting $r = \rho - \overline{(\rho)}^{\frac{1}{\varepsilon}}$:

$$\frac{\partial}{\partial t}(r) + \operatorname{div}(ur) \leq 0. \quad (2.35)$$

We now want to integrate and to use the fact that $r \geq 0$ to get that $r = 0$ a.a. To justify the integration we test our inequality against a cut-off function of the form $\varphi(\frac{\cdot}{R})$ where $\varphi \in C_0^\infty(\mathbb{R}^N)$, $\varphi = 1$ on $B(0, 1)$, $\operatorname{Supp} \varphi \subset B(0, 2)$ and $R > 1$. We test the equation (2.35) against φ_R and we get:

$$\int_{[0, T] \times \mathbb{R}^N} \frac{\partial}{\partial t}[r(t, x)] \varphi_R(x) - u(t, x) r(t, x) \frac{1}{R} \nabla \varphi\left(\frac{x}{R}\right) dt dx \leq 0. \quad (2.36)$$

Next we notice that:

$$\begin{aligned} \left| \int_{[0, T] \times \mathbb{R}^N} u(t, x) r(t, x) \frac{1}{R} \nabla \varphi\left(\frac{x}{R}\right) dt dx \right| &\leq \|u\|_{L^1(0, T; L^{\frac{2N}{N-2}}(\mathbb{R}^N))} \|r\|_{L^1(0, T; L^{\frac{2N}{N+2}}(\mathbb{R}^N))} \\ &\quad * \frac{1}{R} \|\nabla \varphi\|_{L^\infty(\mathbb{R}^N)}. \end{aligned}$$

It implies that:

$$\int_{[0, T] \times \mathbb{R}^N} u(t, x) r(t, x) \frac{1}{R} \nabla \varphi\left(\frac{x}{R}\right) dt dx \rightarrow 0 \quad \text{as } R \rightarrow +\infty.$$

We have then:

$$\int_{[0, T] \times \mathbb{R}^N} \frac{\partial}{\partial t} r(t, x) \varphi_R(x) dt dx = \int_{\mathbb{R}^N} r(T, x) \varphi_R(T, x) dx - \int_{\mathbb{R}^N} r(0, x) \varphi_R(0, x) dx.$$

Now we want to verify that $r(0, \cdot) = 0$ to conclude because. Indeed we will obtain that:

$$\lim_{R \rightarrow +\infty} \int_{\mathbb{R}^N} r(T, x) \varphi_R(T, x) dx \rightarrow \int_{\mathbb{R}^N} r(T, x) dx \leq 0 \quad \text{and } r \geq 0.$$

then $r = 0$.

We know that ρ_n is uniformly bounded in $L^\infty(L^1 \cap L^s(\mathbb{R}^N))$, then ρ_n^ε is relatively compact in $C([0, T]; L^p - w)$ with $1 < p < s$ (where $L^p - w$ denote the space L^p endowed with weak topology). Moreover $(\rho_0^\varepsilon)_n$ converges to ρ_0^ε , we deduce then $r(0) = 0$ a.a.

Now as $r = 0$ we conclude in using the theorem 2.4 ρ_n converges a.a to ρ and that ρ_n converges to ρ in $L^p([0, T] \times B_R)$ for all $p \in [1, q)$ and in $L^{p_1}(0, T, L^{p_2}(B_R))$ for all $p_1 \in [1, +\infty)$, $p_2 \in [1, s)$ and for all $R \in (0, +\infty)$.

Conclusion

We wish now conclude and get the convergence of our theorem in the total space.

We aim at proving here the convergence of ρ_n in $C([0, T], L^p(\mathbb{R}^N)) \cap L^{q'}(\mathbb{R}^N \times (0, T))$ for all $1 \leq p < s$, $1 \leq q' < q$. We have just to show the convergence of ρ_n to ρ in $C([0, T], L^1(\mathbb{R}^N))$. To this end, we introduce $d_n = \sqrt{\rho_n}$ which clearly converges to $\sqrt{\rho}$ in $L^{2p_1}(0, T, L^{2p_2}(B_R)) \cap L^{2p}(B_R \times (0, T))$ to $d = \sqrt{\rho}$ for all $R \in (0, +\infty)$.

We next remark that $\rho \in C([0, T], L^1(\mathbb{R}^N))$ and thus $d \in C([0, T], L^2(\mathbb{R}^N))$. Indeed,

using once more the regularization lemma in [14] we obtain the existence of a bounded $\rho_\alpha \in C([0, T], L^1(\mathbb{R}^N))$ smooth in x for all t satisfying:

$$\frac{\partial \rho_\alpha}{\partial t} + \operatorname{div}(u \rho_\alpha) = r_\alpha \quad \text{in } L^1((0, T) \times \mathbb{R}^N) \quad \text{as } \alpha \rightarrow 0_+.$$

with $r_\alpha = \operatorname{div}(u \rho_\alpha) - \operatorname{div}(\rho u) * w_\alpha$ (where w is defined as in the previous part).
 $\rho_\alpha \rightarrow \rho$ in $L^1(\mathbb{R}^N \times (0, T))$, $\rho_\alpha|_{t=0} \rightarrow \rho|_{t=0}$ in $L^1(\mathbb{R}^N)$ as $\alpha \rightarrow 0_+$.
 From these facts, it is straightforward to deduce that:

$$\frac{\partial}{\partial t} |\rho_\alpha - \rho_\eta| + \operatorname{div}(u |\rho_\alpha - \rho_\eta|) = r_\alpha - r_\eta$$

and thus:

$$\sup_{[0, T]} \int_{\mathbb{R}^N} |\rho_\alpha - \rho_\eta| dx = \int_0^T \int_{\mathbb{R}^N} |r_\alpha - r_\eta| dx.$$

Since $\rho \in C([0, T], L^p(B_R) - w)$ (for all $R \in (0, +\infty)$, $1 < p < s$), we may then deduce that ρ_α converge to ρ in $C([0, T], L^1(\mathbb{R}^N))$.

Next, we observe that we can justify as we did above that d_n and d satisfy:

$$\begin{aligned} \frac{\partial d_n}{\partial t} + \operatorname{div}(u_n d_n) &= \frac{1}{2} d_n \operatorname{div}(u_n), \\ \frac{\partial d}{\partial t} + \operatorname{div}(u d) &= \frac{1}{2} d \operatorname{div}(u). \end{aligned}$$

Therefore once more, d_n converges to d in $C([0, T], L^2(\mathbb{R}^N) - w)$.

Thus in order to conclude, we just have to show that whenever $t_n \in [0, T]$, $t_n \rightarrow t$, then $d_n(t_n) \rightarrow d(t)$ in $L^2(\mathbb{R}^N)$ or equivalently that $\int_{\mathbb{R}^N} d_n(t_n)^2 dx = \int_{\mathbb{R}^N} \rho_n(t_n) dx \rightarrow_n \int_{\mathbb{R}^N} d(t)^2 dx = \int_{\mathbb{R}^N} \rho(t) dx$. This is the case since we deduce from the mass equation, integrating this equation over $\mathbb{R}^N$ and justifying the integration exactly like previously that:

$$\int_{\mathbb{R}^n} \rho_n(t_n) dx = \int_{\mathbb{R}^n} (\rho_0)_n dx \rightarrow_n \int_{\mathbb{R}^n} \rho_0 dx = \int_{\mathbb{R}^n} \rho(t) dx.$$

We then conclude by uniform continuity that $\|\rho_n(t_n) - \rho_n(t)\|_{L^1}$ tends to 0.

Case $N = 2$

First of all, the main difficulty is the fact that we no longer have global L^p bounds on u_n . That's why most of the proof is in fact local and we know that u_n is bounded in $L^2(0, T; L^p(B_R))$ for all $p \in [1, +\infty)$, $R \in (0, +\infty)$.

As we need to localize the argument, we get the following limit:

$$((\mu + \xi) \operatorname{div} u_n - a \rho_n^\gamma - \frac{1}{2} \rho_n^2) \rho_n^\varepsilon \rightarrow_n ((\mu + \xi) \operatorname{div} u - a \overline{\rho}^\gamma - \frac{1}{2} \overline{\rho}^2) \overline{\rho}^\varepsilon \quad \text{in } \mathcal{D}'(\mathbb{R}^N \times [0, T])$$

let $\varphi \in C_0^\infty(\mathbb{R}^N)$, $0 \leq \varphi \leq 1$, $\operatorname{supp} \varphi \subset K$ for an arbitrary compact set $K \in \mathbb{R}^N$. We apply the operator $(\Delta)^{-1} \operatorname{div}$ to the momentum equation that we have localized and we

pass directly to the weak limit:

$$\begin{aligned} \frac{\partial}{\partial t}(-\Delta)^{-1} \operatorname{div}(\varphi \rho u) + R_{ij}(\varphi \rho u_i u_j) + [(\mu + \xi) \operatorname{div} u \varphi - \overline{a \rho^\gamma} \varphi] \\ = \kappa(-\Delta)^{-1} \operatorname{div}(\varphi \overline{\rho(\nabla \phi * \rho)} - \frac{\kappa}{2} \overline{\varphi \rho^2}) + (-\Delta)^{-1} \overline{R}. \end{aligned} \quad (2.37)$$

with:

$$\begin{aligned} \overline{R} &= \partial_i \varphi \partial_j (\rho u_i u_j) + (\partial_{ij} \varphi) \rho u_i u_j - (\mu + \xi) \Delta \varphi \operatorname{div} u - (2\mu + \xi) \nabla \varphi \cdot \nabla \operatorname{div} u + \mu \Delta u \cdot \nabla \varphi \\ &+ \Delta \varphi a \overline{\rho^\gamma} + a \nabla \varphi \cdot \nabla \overline{\rho^\gamma} + \frac{\kappa}{2} \Delta \varphi \overline{\rho^2} + \frac{\kappa}{2} \nabla \varphi \cdot \nabla \overline{\rho^2}. \end{aligned}$$

Now we multiply (2.37) with $\overline{\rho^\varepsilon}$ and we verify that each term has a sense.

So we get in proceeding in the same way as before, we can verify that $(\rho_n)^\varepsilon (-\Delta)^{-1} R_n$ converges in distribution sense to $\overline{\rho^\varepsilon} (-\Delta)^{-1} \overline{R}$ for small enough ε . We get as in the previous case for $N \geq 3$:

$$\varphi[(\mu + \xi) \overline{\operatorname{div} u (\rho)^\varepsilon} - a \overline{\rho^{\gamma+\varepsilon}} - \frac{\kappa}{2} \overline{\rho^{2+\varepsilon}}] = \varphi[(\mu + \xi) (\operatorname{div} u) \varphi \overline{\rho^\varepsilon} - a \overline{\varphi \rho^\gamma} \overline{\rho^\varepsilon} - \frac{\kappa}{2} \overline{\varphi \rho^2} \overline{\rho^\varepsilon}] \quad \text{a.e.}$$

We then deduce the following inequalities as in the previous proof:

$$\frac{d}{dt} (\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}} + \operatorname{div}(u (\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}}) \geq 0 \quad \text{in } \mathcal{D}'((0, T) \times \mathbb{R}^N).$$

We see that the only point left to check is the justification of the integration over $\mathbb{R}^2$ of terms like $\operatorname{div}(\overline{\rho^\varepsilon}^{\frac{1}{\varepsilon}} u)$ or $\operatorname{div}(\rho u)$ and more precisely that the integral vanishes. This is in fact straightforward provided we use the bounds on $\rho \in L^\infty(L^1(\mathbb{R}^N))$ and $\rho |u|^2 \in L^\infty(L^1(\mathbb{R}^N))$ and so $\rho u \in L^\infty(L^1(\mathbb{R}^N))$. Then, letting $\varphi \in C_0^\infty(\mathbb{R}^2)$, $0 \leq \varphi \leq 1$, $\varphi = 1$ on $B(0, 1)$ and $\varphi = 0$ on ${}^c B(0, 2)$. We set $\varphi_R(\cdot) = \varphi(\frac{\cdot}{R})$ for $R \geq 1$, we have similarly as in the previous case:

$$|\int_0^T dt \int_{\mathbb{R}^2} r u \cdot \nabla \varphi_R(x) dx| \leq \|\nabla \varphi\|_{L^\infty(\mathbb{R}^2)} \frac{1}{R} \|\rho u\|_{L^1(0, T) \times \mathbb{R}^2} \rightarrow 0 \quad \text{as } R \rightarrow +\infty$$

We can then conclude as in the previous proof.

□

Proof of the convergence assertion on $\rho_n u_n$

We now want to show the convergence of $\rho_n u_n$ to have informations on strong convergence of u_n modulo the vacuum. We recall in this part some classical inequalities to get the convergence of $\rho_n u_n$, for more details see Lions in [15]. We use once more a mollifier $k_\alpha = \frac{1}{\alpha^N} k(\frac{\cdot}{\alpha})$ where $k \in C_0^{+\infty}(\mathbb{R}^N)$ and we let $g_\alpha = g * k_\alpha$ for an arbitrary function g . We first observe that we have for all $\frac{N}{2} < p < s$:

$$\begin{aligned} |((\rho_n u_n)_\alpha - \rho_n u_n)(x)| &= |\int_{\mathbb{R}^d} [\rho_n(t, y) - \rho_n(t, x)] u_n(t, y) k_\alpha(x - y) dy \\ &\quad + \rho_n(t, x) ((u_n)_\alpha - u_n)(t, x)| \end{aligned}$$

We have in using Hölder inequalities with the measure $k_\alpha(x-y)dy$:

$$|((\rho_n u_n)_\alpha - \rho_n u_n)(x)| \leq \left[\int_{\mathbb{R}^d} |\rho_n(t, y) - \rho_n(t, x)|^p k_\alpha(x-y) dy \right]^{\frac{1}{p}} \left(|u_n|^{\frac{p}{p-1}} \right)_\alpha^{\frac{p-1}{p}} + \rho_n |(u_n)_\alpha - u_n|(t, x).$$

Hence for all $t \geq 0$

$$\begin{aligned} \int_{\mathbb{R}^d} |((\rho_n u_n)_\alpha - \rho_n u_n)(x)| dx &\leq \left[\int_{\mathbb{R}^d} dx \int_{\mathbb{R}^d} |\rho_n(t, y) - \rho_n(t, x)|^p k_\alpha(x-y) dy \right]^{\frac{1}{p}} \\ &\quad * \left\| \left(|u_n|^{\frac{p}{p-1}} \right)_\alpha \right\|_{L^1}^{\frac{p-1}{p}} + \|\rho_n\|_{L^p} \|(u_n)_\alpha - u_n\|_{L^{\frac{p-1}{p}}}, \\ &\leq \left[\sup_{|z| \leq \alpha} \|\rho_n(\cdot + z) - \rho_n\|_{L^p} \right] \|u_n\|_{L^{\frac{p-1}{p}}} + \|\rho_n\|_{L^p} \|(u_n)_\alpha - u_n\|_{L^{\frac{p-1}{p}}}. \end{aligned}$$

Next if we choose $p > \frac{2N}{N+2}$, $\frac{p}{p-1} < \frac{2N}{N-2}$ then $\|(u_n)_\alpha - u_n\|_{L^2(0,T;L^{\frac{p-1}{p}})}$ converge to 0 as α goes to 0_+ uniformly in n . In addition, the convergence on ρ_n assure that $\sup_{|z| \leq \alpha} \|\rho_n(\cdot + z) - \rho_n\|_{L^p}$ converge to 0 as α goes to 0_+ uniformly in n . Therefore in conclusion, $(\rho_n u_n)_\alpha - \rho_n u_n$ converge to 0 in $L^2(0, T; L^1)$ as α goes to 0_+ uniformly in n . Next $(\rho_n u_n)_\alpha$ is smooth in x , uniformly in n and in $t \in [0, T]$. Therefore, remarking that $\frac{\partial}{\partial t}(\rho_n u_n)_\alpha$ is bounded in a $L^2(0, T; H^m)$ for any $m \geq 0$, we deduce that $(\rho_n u_n)_\alpha$ converge to $(\rho u)_\alpha$ as n goes to $+\infty$ in $L^1((0, T) \times \mathbb{R}^N)$ for each α . Then using the bound on $\rho_n u_n$ in $L^\infty(L^{\frac{2s}{s+1}})$, we deduce that $\rho_n u_n$ converge to ρu in $L^1((0, T) \times \mathbb{R}^N)$ and we can conclude by interpolation.

The last convergence is a consequence of the strong convergence of ρ_n and $\rho_n u_n$.

3 Existence of weak solution with general pressure

In the sequel we concentrate us only on the cases $N = 2, 3$. We now want to extend our previous result to more general and physical pressure laws. In particular we are now interested by two cases, the first one concerns monotonous pressure law (close in a certain sense that we will precise to ρ^γ pressure), the second one is the case of a slightly modified Van der Waals pressure.

The technics of proof will be very similar to the previous proof, only technical points change.

3.1 Monotonous pressure

In this section, we shall investigate an extension of the preceding results to the case of a general monotonous pressure P , i.e P is assumed to be a C^1 non-decreasing function on $[0, +\infty)$ vanishing at 0.

We want here to mention in the general situation our new energy inequality, we recall the inequality (1.4):

$$\begin{aligned} \int_{\mathbb{R}^N} \left(\frac{1}{2} \rho |u|^2 + \Pi(\rho) + E_{global}[\rho(\cdot, t)] \right)(x) dx(t) + \int_0^t \int_{\mathbb{R}^N} (\mu D(u) : D(u) \\ + (\lambda + \mu) |\operatorname{div} u|^2) dx \leq \int_{\mathbb{R}^N} \left(\frac{|m_0|^2}{2\rho} + \Pi(\rho_0) + E_{global}[\rho_0] \right) dx. \end{aligned}$$

where we define Π by: $\frac{\partial}{\partial t}(\frac{\Pi(t)}{t}) = \frac{P(t)}{t^2}$ for all $t > 0$.

There are two cases worth considering: first of all if $P(t)$ is such that $\int_0^1 \frac{P(s)}{s^2} ds < +\infty$ then we can choose $\Pi(\rho) = \rho \int_0^\rho \frac{P(s)}{s^2} ds$.

In the other case, i.e $\lim_{t \rightarrow 0} \frac{P(t)}{t} = c > 0$, we can choose $\Pi(\rho) = \rho \int_\rho^1 \frac{P(s)}{s^2} ds$ and Π behaves like $\rho \log \rho$ as ρ goes to 0.

We now consider a sequence of solutions (ρ_n, u_n) and we make the same assumptions on this sequence as in the previous section except that we need to modify the assumptions on ρ_n . We assume always that ρ_n is bounded in $C([0, T], L^1(\mathbb{R}^N))$, $P(\rho_n)$ is bounded in $L^\infty(0, T, L^1(\mathbb{R}^N))$.

$(\rho_n)_{n \geq 1}$ is bounded in $L^\infty(0, T, L^2)$,

and we also assume that we have:

$$(\rho_n^\varepsilon P(\rho_n))_{n \geq 1} \text{ is bounded in } L^1(K \times (0, T))$$

for some $\varepsilon > 0$, where K is an arbitrary compact set included in $\mathbb{R}^N$.

Theorem 3.6 *Let the assumptions of theorem 1.1 be satisfied with in addition P a monotone pressure.*

Then there exists a renormalized finite energy weak solution to problem (NSK) in the sense of definitions 1.1 and 1.2. Moreover $P(\rho_n)$ converges to $P(\rho)$ in $L^1(K \times (0, T))$ for any compact set K .

Proof:

In this situation, the proof of theorem 3.6 is based on the same compactness argument as in the theorem 1.1. In particular, there is essentially one observation which allows us to adapt the proof of theorem 1.1. Namely we still obtain the following identity for the effective viscous flux:

$$\beta(\rho_n)((\mu + \xi)\operatorname{div} u_n - P(\rho_n) - \frac{\kappa}{2}\rho_n^2) \rightharpoonup_n \overline{\beta(\rho)}((\mu + \xi)\operatorname{div} u - \overline{P(\rho)} - \frac{\kappa}{2}\overline{\rho^2}),$$

with $\beta(\rho) = \rho^\varepsilon$ for ε small enough. We have then:

$$(\mu + \xi)\overline{\rho^\varepsilon \operatorname{div} u} - \overline{P(\rho)\rho^\varepsilon} - \frac{\kappa}{2}\overline{\rho^{2+\varepsilon}} = (\mu + \xi)\overline{\operatorname{div} u \rho^\varepsilon} - \overline{P(\rho)\rho^\varepsilon} - \frac{\kappa}{2}\overline{\rho^2 \rho^\varepsilon}, \quad (3.38)$$

Now we can recall a lemma coming from P.-L.Lions in [15]:

Lemma 2 *Let $p_1, p_2 \in C([0, \infty))$ be non-decreasing functions. We assume that $p_1(\rho_n), p_2(\rho_n)$ are relatively weakly compact in $L^1(K \times (0, T))$ for any compact set $K \subset \mathbb{R}^N$. Then, we have:*

$$\overline{p_1(\rho)p_2(\rho)} \geq \overline{p_1(\rho)p_2(\rho)} \quad a.e.$$

We get finally as in the proof of theorem 1.1 in using lemma 2:

$$\overline{\operatorname{div} u \rho^\varepsilon} \geq \operatorname{div} u \overline{\rho^\varepsilon},$$

All remaining argumentation of the proof of theorem 1.1 can be performed to conclude.

□

3.2 Pressure of Van der Waals type

In this section we are interested by pressure of type Van der Waals which consequently are not necessarily non-decreasing. That's why in the following proof we will proceed slightly differently.

So we consider the pressure law:

$$P(\rho) = \frac{RT_*\rho}{b-\rho} - a\rho^2 \text{ for } \rho \leq b - \theta \text{ for some small } \theta > 0$$

and we extend the function P to be strictly increasing function on $\rho \geq b - \theta$.

We have then:

1. P' is bounded from below, that is:

$$P'(\rho) \geq -\bar{\rho} \text{ for all } \rho > 0.$$

2. P is a strictly increasing function for ρ large enough.

Under the above conditions, it is easy to see that the pressure can be written as:

$$P(\rho) = P_1(\rho) - P_2(\rho).$$

with P_1 a non-decreasing function of ρ , and

$$P_2 \in C^2[0, +\infty), \quad P_2 \geq 0, \quad P_2 = 0 \text{ for } \rho \geq \bar{\rho}.$$

Remark 1 *The a priori energy estimate give us the bound of ρ in $L^\infty(L^2)$, then we have:*

$$|\{(t, x) \in (0, T) \times \mathbb{R}^N / |\rho(t, x)| > b\}| \leq \frac{T \|\rho\|_{L^\infty(L^2(\mathbb{R}^N))}^2}{b^2}.$$

Then we can controll in measure the set where P is different from the Van der Waals measure, and it's a set of finite measure.

We obtain the following theorem.

Theorem 3.7 *If in addition to the above assumptions, we assume that ρ_n^0 converges in $L^1(\mathbb{R}^N)$ to ρ_0 then (ρ, u) is a weak solution of the system (NSK) satisfying the initial condition and we have:*

$$\rho_n \rightarrow \rho \text{ in } C([0, T], L^p(\mathbb{R}^N)) \cap L^r((0, T) \times \mathbb{R}^N) \text{ for all } 1 \leq p < 2, 1 \leq r < 1 + \frac{4}{N},$$

Proof:

Most of the proof of theorem 3.7 is similar as theorem 1.1. We will use a approximated sequel T_k (introduced by Feireisl in [?]) of ρ by some concave bounded function.

Definition 3.3 We define the function $T \in C^\infty(\mathbb{R}^N)$ as follows:

$$\begin{aligned} T(z) &= z \quad \text{for } z \in [0, 1], \\ T(z) &\text{ concave on } [0, +\infty), \\ T(z) &= 2 \quad \text{for } z \geq 3, \\ T(z) &= -T(-z) \quad \text{for } z \in (-\infty, 0], \end{aligned}$$

And T_k is the cut-off function:

$$T_k(z) = kT\left(\frac{z}{k}\right).$$

In following the proof of theorem 1.1, we get:

$$\begin{aligned} \frac{\partial}{\partial t}(\overline{L_k(\rho)} - L_k(\rho)) + \operatorname{div}((\overline{T_k(\rho)} - T_k(\rho))u) + \overline{T_k(\rho)}\operatorname{div}u - \overline{T_k(\rho)}\operatorname{div}u \\ = (T_k(\rho) - \overline{T_k(\rho)})\operatorname{div}u \quad \text{in } \mathcal{D}'((0, T) \times \mathbb{R}^N), \end{aligned}$$

where $L_k(\rho) = \rho \log(\rho)$ for $0 \leq \rho \leq k$, and $0 \leq L_k(\rho) \leq \rho \log(\rho)$ otherwise.

So we get in integrating in time on $[t_1, t_2]$:

$$\begin{aligned} \int_{\mathbb{R}^N} (\overline{L_k(\rho)} - L_k(\rho))(t_2)dx - \int_{\mathbb{R}^N} (\overline{L_k(\rho)} - L_k(\rho))(t_1)dx \\ + \int_{t_1}^{t_2} \int_{\mathbb{R}^N} \overline{T_k(\rho)}\operatorname{div}u - \overline{T_k(\rho)}\operatorname{div}u \, dxdt = \int_{t_1}^{t_2} \int_{\mathbb{R}^N} (T_k(\rho) - \overline{T_k(\rho)})\operatorname{div}u \, dxdt. \end{aligned}$$

We can show that:

$$\|T_k(\rho) - \overline{T_k(\rho)}\|_{L^2((0, T) \times \mathbb{R}^N)} \rightarrow_{k \rightarrow +\infty} 0$$

For proving the previous inequality, we see that $\|T_k(\rho) - \overline{T_k(\rho)}\|_{L^1((0, T) \times \mathbb{R}^N)} \rightarrow 0$ for $k \rightarrow +\infty$. We then conclude by interpolation with $T_k(\rho) - \overline{T_k(\rho)} \in L^q(0, T) \times \mathbb{R}^N$ with $q > 2$. By Hölder inequality we obtain that:

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^N} (T_k(\rho) - \overline{T_k(\rho)})\operatorname{div}u \, dxdt \rightarrow 0 \quad \text{for } k \rightarrow +\infty.$$

We have then:

$$\begin{aligned} \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^N} (\overline{L_k(\rho)} - L_k(\rho))(t_2)dx - \int_{\mathbb{R}^N} (\overline{L_k(\rho)} - L_k(\rho))(t_1)dx \\ = - \lim_{k \rightarrow +\infty} \int_{t_1}^{t_2} \int_{\mathbb{R}^N} \overline{T_k(\rho)}\operatorname{div}u - \overline{T_k(\rho)}\operatorname{div}u \, dxdt. \end{aligned} \tag{3.39}$$

We set:

$$\begin{aligned} df t[\rho_n \rightarrow \rho](t) &= \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^N} (\overline{L_k(\rho)} - L_k(\rho))(t)dx \\ A(k, \rho) &= \int_{t_1}^{t_2} \int_{\mathbb{R}^N} \overline{T_k(\rho)}\operatorname{div}u - \overline{T_k(\rho)}\operatorname{div}u \, dxdt. \end{aligned}$$

We can show as in the previous proof of theorem 1.1 that:

$$\begin{aligned} \int_{t_1}^{t_2} \int_K \overline{T_k(\rho) \operatorname{div} u} - \overline{T_k(\rho)} \operatorname{div} u \, dx dt \\ = \lim_{n \rightarrow +\infty} \int_{t_1}^{t_2} \int_K \left((P(\rho_n) + \frac{\kappa}{2} \rho_n^2) T_k(\rho_n) - \overline{(P(\rho) + \frac{\kappa}{2} \rho^2) T_k(\rho)} \right) dx dt. \end{aligned}$$

for any compact $K \subset \mathbb{R}^N$.

Using the lemma 2 we deduce that:

$$\lim_{n \rightarrow +\infty} \int_{t_1}^{t_2} \int_{\mathbb{R}^N} \left(P_1(\rho_n) + \frac{\kappa}{2} \rho_n^2 \right) T_k(\rho_n) - \overline{(P_1(\rho) + \frac{\kappa}{2} \rho^2) T_k(\rho)} \, dx dt \leq 0.$$

We have then:

$$\begin{aligned} dft[\rho_n \rightarrow \rho](t_2) - dft[\rho_n \rightarrow \rho](t_1) \leq \lim_{k \rightarrow +\infty} \left(\lim_{n \rightarrow +\infty} \int_{t_1}^{t_2} \int_{\mathbb{R}^N} P_2(\rho_n) T_k(\rho_n) \right. \\ \left. - \overline{P_2(\rho) T_k(\rho)} \, dx dt \right). \end{aligned}$$

As the sequence $(\rho_n)_{n \in \mathbb{N}}$ is bounded in $L^\infty(0, T, L^2(\mathbb{R}^N))$, and P_2 is a bounded function, we have:

$$\begin{aligned} \lim_{k \rightarrow +\infty} \left(\lim_{n \rightarrow +\infty} \int_{t_1}^{t_2} \int_{\mathbb{R}^N} P_2(\rho_n) T_k(\rho_n) - \overline{P_2(\rho) T_k(\rho)} \, dx dt \right) \\ = \lim_{n \rightarrow +\infty} \int_{t_1}^{t_2} \int_{\mathbb{R}^N} P_2(\rho_n) \rho_n - \overline{P_2(\rho)} \bar{\rho} \, dx dt. \end{aligned}$$

Since the function P_2 is twice continuously differentiable and compactly supported in $[0, +\infty)$, there exists $\Lambda > 0$ big enough such that both $\rho \rightarrow \Lambda \rho \log \rho - \rho P_2(\rho)$ and $\rho \rightarrow \Lambda \rho \log \rho + \rho P_2(\rho)$ are convex functions of ρ , indeed the twice derivative are positive. As a consequence of weak lower semi-continuity of convex functionals, we obtain:

$$\begin{aligned} \lim_{n \rightarrow +\infty} \int_{t_1}^{\tau_2} \int_{\mathbb{R}^N} P_2(\rho_n) \rho_n - \overline{P_2(\rho)} \bar{\rho} \, dx dt \\ \leq \Lambda \int_{t_1}^{\tau_2} \int_{\mathbb{R}^N} (\overline{\rho \log \rho} - \rho \log \rho) \, dx dt + \int_{t_1}^{\tau_2} \int_{\mathbb{R}^N} (P_2(\rho) - \overline{P_2(\rho)}) \rho \, dx dt. \end{aligned}$$

Futhermore we have:

$$\begin{aligned} \int_{t_1}^{t_2} \int_{\mathbb{R}^N} (P_2(\rho) - \overline{P_2(\rho)}) \rho \, dx dt &\leq \int_{t_1}^{\tau_2} \int_{\rho \leq \rho_r} (P_2(\rho) - \overline{P_2(\rho)}) \rho \\ &\leq \Lambda \int_{t_1}^{t_2} \int_{\rho \leq \rho_r} (\overline{\rho \log \rho} - \rho \log \rho) \, dx dt \\ &\leq \Lambda \rho_r \int_{t_1}^{t_2} \int_{\mathbb{R}^N} (\overline{\rho \log \rho} - \rho \log \rho) \, dx dt \end{aligned}$$

The previous relation gives:

$$dft[\rho_n \rightarrow \rho](t_2) \leq dft[\rho_n \rightarrow \rho](t) + \omega \int_{t_1}^{t_2} dft[\rho_n \rightarrow \rho](t).$$

Applying Grönwall's lemma we infer:

$$dft[\rho_n \rightarrow \rho](t_2) \leq dft[\rho_n \rightarrow \rho](t_1) \exp(\omega(t_2 - t_1)).$$

We conclude that $dft[\rho_n \rightarrow \rho](t) = 0 \forall t$, because ρ_n^0 converges strongly in L^1 to ρ_0 .

4 Weak solutions with data close to a stable equilibrium

We consider in this section one situation which is rather different from the three cases considered in the preceeding sections. This situation is relevant for practical applications and realistic flow and they involve conditions at infinity different from those studied.

We wish to investigate the system (NSK) with hypothesis close from these of the theorem for strong solutions. We want then to study the system with a density close from a stable equilibrium in the goal to can choose initial data which avoid the vacuum. We look now for a solution (ρ, u) defined on $\mathbb{R} \times \mathbb{R}^N$ of the system (NSK) (where $P(\rho) = a\rho^\gamma$) with $\rho \geq 0$ on $\mathbb{R} \times \mathbb{R}^N$.

In addition we require (ρ, u) to satisfy the following limit conditions:

$$(\rho, u)(x, t) \rightarrow (\bar{\rho}, 0) \text{ as } |x| \rightarrow +\infty, \text{ for all } t > 0$$

where $\bar{\rho} > 0$.

Such an analysis requires the use of the Orlicz spaces. We define the Orlicz space $L_p^q(\mathbb{R}^N)$ as follows:

$$L_p^q(\mathbb{R}^N) = \{f \in L_{loc}^1(\mathbb{R}^N) / f1_{\{|f| \leq \delta\}} \in L^p(\mathbb{R}^N), f1_{\{|f| \geq \delta\}} \in L^q(\mathbb{R}^N)\}.$$

For the following properties, we can give this definition too.

Definition 4.4 *We define Ψ as a convex function on $[0, +\infty)$ which is equal (or equivalent) to x^p for x small and to x^q for x large.*

$$L_p^q(\mathbb{R}^N) = \{f \in L_{loc}^1(\mathbb{R}^N) / \Psi(f) \in L^1(\mathbb{R}^N)\}.$$

We can check that $L_p^q(\mathbb{R}^N)$ is a linear space. Now we endow $L_p^q(\mathbb{R}^N)$ with a norm so that $L_p^q(\mathbb{R}^N)$ is a separable Banach space.

$$\|f\|_{L_p^q(\mathbb{R}^N)} = \inf \left(t > 0 / \Psi\left(\frac{f}{t}\right) \leq 1 \right).$$

We recall now some useful properties of the Orlicz space:

Corollary 1 *We have:*

1. *Embedding:*

$$L_p^q(\mathbb{R}^N) \subset L_{p_1}^{q_1}(\mathbb{R}^N) \text{ if } 1 \leq q_1 \leq q < +\infty, 1 \leq p \leq p_1 < +\infty.$$

2. *Topology: $f_n \rightarrow_n 0$ in $L_p^q(\mathbb{R}^N)$ if and only if $\Psi(f_n) \rightarrow_n 0$ and that:*

$$\Psi\left(\frac{f}{\|f\|_{L_p^q(\mathbb{R}^N)}}\right) = 1 \text{ if } f \neq 0$$

We recall now some useful properties of the Orlicz space:

Proposition 4.1 *We then have:*

- *Dual space: If $p > 1$ then $(L_p^q(\mathbb{R}^N))' = L_{p'}^{q'}(\mathbb{R}^N)$ where $q' = \frac{q}{q-1}$, $p' = \frac{p}{p-1}$.*
- *Let F be a continuous function on $\mathbb{R}$ such that $F(0) = 0$, F is differentiable at 0 and $F(t)|t|^{-\theta} \rightarrow \alpha \neq 0$ at $t \rightarrow +\infty$. Then if $q \geq \theta$,*

$$F(f) \in L_{\frac{q}{\theta}}^{\frac{q}{\theta}}(\mathbb{R}^N) \text{ if } f \in L_p^q(\mathbb{R}^N).$$

Our goal is now to get energy estimate. We have to face a new difficulty. Indeed ρ , $\rho|u|^2$, ρ^γ need not belong to L^1 .

We first want to explain how it is possible to obtain natural a priori bounds which correspond to energy-like identities.

Next we write the following formal identities:

$$\begin{aligned} & \frac{1}{\gamma-1} \frac{d}{dt} (\rho^\gamma - \bar{\rho}^\gamma - \gamma \bar{\rho}^{\gamma-1} (\rho - \bar{\rho}) + \operatorname{div} [u \frac{\gamma}{\gamma-1} (\rho^\gamma - \bar{\rho}^{\gamma-1} \rho)]) = u \cdot \nabla (\rho^\gamma) \\ & \rho \frac{d}{dt} \frac{|u|^2}{2} + \rho u \cdot \nabla \frac{|u|^2}{2} - \mu \Delta u \cdot u - \xi \nabla \operatorname{div} u \cdot u + a u \cdot \nabla \rho^\gamma = \kappa \rho u \nabla (\phi * \rho - \rho). \end{aligned} \quad (4.40)$$

We may then integrate in space the equality (4.40) and we get:

$$\begin{aligned} & \left(\int_{\mathbb{R}^N} \rho \frac{|u|^2}{2} + \frac{a}{\gamma-1} (\rho^\gamma + (\gamma-1) \bar{\rho}^\gamma - \gamma \bar{\rho}^{\gamma-1} \rho) + E_{\text{global}}[\rho - \bar{\rho}] dx \right) (t) \\ & + \int_0^t ds \int_{\mathbb{R}^N} 2\mu |Du|^2 + 2\xi |\operatorname{div} u|^2 dx \leq \int_{\mathbb{R}^N} \rho_0 \frac{|u_0|^2}{2} + \frac{a}{\gamma-1} (\rho_0^\gamma + (\gamma-1) \bar{\rho}^\gamma - \gamma \bar{\rho}^{\gamma-1} \rho_0) \\ & + E_{\text{global}}[\rho_0 - \bar{\rho}] dx. \end{aligned} \quad (4.41)$$

Notation 2 *In the sequel we will note:*

$$j_\gamma(\rho) = \rho^\gamma + (\gamma-1) \bar{\rho}^\gamma - \gamma \bar{\rho}^{\gamma-1} \rho.$$

Now we can recall a theorem (see [15]) on the Orlicz space concerning this quantity:

Theorem 4.8 *$j_\gamma(\rho) \in L^1(\mathbb{R}^N)$ if and only if $\rho - \bar{\rho} \in L_2^\gamma$.*

In following this theorem and our energy estimate we get that $\rho - \bar{\rho} \in L^\infty(0, T; L_2^\gamma(\mathbb{R}^N))$ for all $T \in \mathbb{R}$.

Moreover we have:

$$E_{\text{global}}[\rho(t, \cdot) - \bar{\rho}](x) = \frac{\kappa}{4} (\rho - \bar{\rho})^2 + \phi * (\rho - \bar{\rho})^2 - 2(\rho - \bar{\rho}) (\phi * (\rho - \bar{\rho})).$$

Then in using the fact that $\rho - \bar{\rho} \in L^\infty(0, T; L_2^\gamma(\mathbb{R}^N)) \hookrightarrow L^2(\mathbb{R}^N) + L^\gamma(\mathbb{R}^N)$ and interpolation on $\nabla \phi$, we get that $\rho - \bar{\rho} \in L^\infty(0, T; L^2(\mathbb{R}^N))$.

We may now turn to our compactness result. First of all, we consider sequences of solutions (ρ_n, u_n) of the system corresponding to some initial conditions (ρ_n^0, u_n^0) . In using the above energy inequalities, we assume that $j_\gamma(\rho_n^0)$, $E_{global}[\rho_n^0 - \bar{\rho}]$ and $\rho_n^0 |u_n^0|^2$ are bounded in $L^\infty(L^1(\mathbb{R}^N))$ and that $\rho_n^0 - \bar{\rho}$ converges weakly in $L_2^\gamma(\mathbb{R}^N)$ to some $\rho_0 - \bar{\rho}$.

We now assume that:

$$j_\gamma(\rho_n), E_{global}[\rho_n - \bar{\rho}], \rho_n |u_n - \bar{u}|^2 \text{ are bounded in } L^\infty(0, T, L^1(\mathbb{R}^N)),$$

Moreover we have for all $T \in (0, +\infty)$ and for all compact sets $K \subset \mathbb{R}^N$:

$$\rho_n - \bar{\rho} \in L^\infty(L^2(\mathbb{R}^N)) \text{ and } \rho_n \text{ is bounded in } L^q((0, T) \times K),$$

for some $q > s$.

$$Du_n \text{ is bounded in } L^2(\mathbb{R}^N \times (0, T))$$

$$u_n \text{ is bounded in } L^2(0, T, H^1(B_R)) \text{ for all } R, T \in (0, +\infty).$$

Extracting subsequences if necessary, we may assume that ρ_n, u_n converge weakly respectively in $L^2((0, T) \times B_R)$, $L^2(0, T; H^1(B_R))$ to ρ, u for all $R, T \in (0, +\infty)$. We also extract subsequences for which $\rho_n u_n, \rho_n u_n \otimes u_n$ converge weakly as previously.

Remark 2 We notice that the conditions at infinity are implicitly contained in the fact that $(\rho_n - \bar{\rho})^2$ and $\rho_n |u_n|^2 \in L^1(\mathbb{R}^N)$.

We then have the following theorem.

Theorem 4.9 Let $\gamma \geq 1$. We assume that ρ_n^0 converges in $L^1(B_R)$ (for all $R \in (0, +\infty)$) to ρ_0 . Then $(\rho_n, u_n)_{n \in \mathbb{N}}$ converges in distribution sense to (ρ, u) a solution of (NSK). Moreover we have for all $R, T \in (0, +\infty)$:

$$\rho_n \rightarrow_n \rho \text{ in } C([0, T], L^p(B_R \times (0, T))) \cap L^{s_1}(B_R \times (0, T)) \text{ for all } 1 \leq p < s, 1 \leq s_1 < q.$$

$$\text{with } q = s + \frac{4}{N} - 1.$$

Proof:

As in the theorem 1.1, we want test the strong convergence of ρ_n on concav function B . Since the proof is purely local, we have again for small enough $\varepsilon > 0$:

$$(\rho_n)^\varepsilon ((\mu + \xi) \operatorname{div} u_n - a(\rho_n)^\gamma - \frac{\kappa}{2} \rho_n^2) \rightharpoonup_n \overline{(\rho)^\varepsilon} ((\mu + \xi) \operatorname{div} u - a \overline{\rho}^\gamma - \frac{\kappa}{2} \overline{\rho}^2) \\ \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}^N),$$

$$\frac{d}{dt}(\overline{\rho^\varepsilon}) + \operatorname{div}(u \overline{\rho^\varepsilon}) \geq (1 - \varepsilon)(\operatorname{div} u) \overline{\rho^\varepsilon} \text{ in } \mathcal{D}'((0, \infty) \times \mathbb{R}^N). \quad (4.42)$$

Next since $(\overline{\rho^\varepsilon})^{\frac{1}{\varepsilon}} \in L^2(B_R \times (0, T))$ for all $R, T \in (0, +\infty)$, as in the theorem 1.1 in using a result of type Diperna-Lions on renormalized solutions, we get:

$$\frac{d}{dt}(\overline{(\rho)^\varepsilon})^{\frac{1}{\varepsilon}} + \operatorname{div}(u \overline{(\rho)^\varepsilon})^{\frac{1}{\varepsilon}} \geq 0 \text{ in } \mathcal{D}'((0, \infty) \times \mathbb{R}^N). \quad (4.43)$$

while we have $\overline{(\rho)^\varepsilon}^{\frac{1}{\varepsilon}} \leq \rho$ a.e in $\mathbb{R}^N \times (0, +\infty)$ and $\overline{(\rho)^\varepsilon}^{\frac{1}{\varepsilon}}_{t=0} = \rho_{t=0}$ in $\mathbb{R}^N$.

Now in subtracting the second equality of (4.43) from the mass equation and setting $f = \rho - \overline{\rho^\varepsilon}^{\frac{1}{\varepsilon}}$, we have:

$$\frac{d}{dt}(f) + \operatorname{div}(uf) \leq 0, \quad f \geq 0 \quad \text{a.e.}, \quad f|_{t=0} = 0 \quad \text{in } \mathbb{R}^N. \quad (4.44)$$

Next we want again to show from (4.44) that $f = 0$, in integrating (4.44) and in using the fact that $f \leq 0$ to conclude that $f = 0$. The difference with the proof of theorem 1.1 is to justify the integration by parts as we work in different energy space.

We need a cut-off function. We introduce $\varphi \in C_0^\infty(\mathbb{R}^N)$, $0 \leq \varphi \leq 1$, $\operatorname{supp} \varphi \subset B_2$, $\varphi = 1$ on B_1 and we set $\varphi_R = \varphi(\frac{x}{R})$ for $R \geq 1$. Multiplying (4.44) by $\varphi_R(x)$, we obtain:

$$\frac{d}{dt} \int_{\mathbb{R}^N} f \varphi_R(x) dx = \int_{\mathbb{R}^N} \frac{1}{R} f u \cdot \nabla \varphi\left(\frac{x}{R}\right). \quad (4.45)$$

Next, if $T > 0$ is fixed, we see that $\operatorname{supp} \nabla \varphi(\frac{\cdot}{R}) \subset \{R \leq |x| \leq 2R\}$, therefore, for R large enough we have:

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^N} f \varphi_R(x) dx &= \int_{\mathbb{R}^N} \frac{1}{R} f u \cdot \nabla \varphi\left(\frac{x}{R}\right), \\ &\leq \frac{C}{R} \int_{\mathbb{R}^N} f |u| 1_{(R \leq |x| \leq 2R)} dx, \quad \text{for } t \in (0, T), \end{aligned} \quad (4.46)$$

To conclude that $f = 0$, we only have to prove that:

$$\frac{1}{R} \int_{\mathbb{R}^N} f |u| 1_{(R \leq |x| \leq 2R)} dx \rightarrow 0 \quad \text{as } R \rightarrow +\infty. \quad (4.47)$$

We now use that fact that $f \in L^\infty(0, T, L^2(\mathbb{R}^N))$ and $f|u|^2 \in L^\infty(0, T, L^1(\mathbb{R}^N))$ for all $T \in (0, +\infty)$ to control (4.47). The second fact is obvious since $0 \leq \rho$ and $\rho|u|^2 \in L^\infty(0, T; L^1(\mathbb{R}^N))$.

In order to prove the first claim, we only have to show that $\overline{(\rho)^\varepsilon}^{\frac{1}{\varepsilon}} - \bar{\rho} \in L^\infty(0, T, L^2(\mathbb{R}^N))$. We rewrite $(\rho_n)^\varepsilon - (\bar{\rho})^\varepsilon = (\bar{\rho} + (\rho_n - \bar{\rho}))^\varepsilon - (\bar{\rho})^\varepsilon$ is bounded in $L^\infty(0, T, L^2(\mathbb{R}^N))$ in using proposition 4.1 with $F(x) = (\bar{\rho} + x)^\varepsilon - (\bar{\rho})^\varepsilon$. So we have $\sqrt{f} \in L^\infty(L^4(\mathbb{R}^N))$ and we get:

$$\frac{1}{R} \int_0^T dt \int_{\mathbb{R}^N} f |u| 1_{(R \leq |x| \leq 2R)} dx \leq \frac{C_0}{R} \operatorname{meas}(C(0, R, 2R))^{\frac{1}{4}}.$$

We recall that:

$$\operatorname{meas}(C(0, R, 2R)) \sim_{R \rightarrow +\infty} C(n) R^N$$

Then we get:

$$\frac{d}{dt} \int_{\mathbb{R}^N} f \varphi_R(x) dx \rightarrow_{R \rightarrow +\infty} 0$$

and we conclude as in the proof of theorem 1.1.

At this stage, it only remains to show that, for instance, ρ_n converge to ρ in $C([0, T], L^1(B_R))$ for all $R, T \in (0, +\infty)$. In order to do so, we just have to localize the corresponding argument in the proof of theorem 1.1.

Therefore we choose for $R, T \in (0, +\infty)$ fixed, $\varphi \in C_0^\infty(\mathbb{R}^N)$ such that $\varphi = 1$ on B_R , $0 \leq \varphi$ on $\mathbb{R}^N$. Then, we observe that we have:

$$\begin{aligned}\frac{\partial}{\partial t}(\varphi^2 \rho_n) + \operatorname{div}(u_n(\varphi^2 \rho_n)) &= \rho_n u_n \cdot \nabla \varphi^2, \quad \frac{\partial}{\partial t}(\varphi^2 \rho) + \operatorname{div}(u(\varphi^2 \rho)) = \rho u \cdot \nabla \varphi^2 \\ \frac{\partial}{\partial t}(\varphi \sqrt{\rho_n}) + \operatorname{div}(u_n(\varphi \rho_n)) &= \frac{1}{2}(\operatorname{div} u_n) \varphi \sqrt{\rho_n} + \sqrt{\rho_n} u_n \cdot \nabla \varphi, \\ \frac{\partial}{\partial t}(\varphi \sqrt{\rho}) + \operatorname{div}(u(\varphi \rho)) &= \frac{1}{2}(\operatorname{div} u) \varphi \sqrt{\rho} + \sqrt{\rho} u \cdot \nabla \varphi.\end{aligned}$$

From these equations, we deduce as in the proof of the previous theorem, that $\varphi^2 \rho \in C([0, +\infty), L^1(\mathbb{R}^N))$, $\varphi \sqrt{\rho} \in C([0, +\infty), L^2(\mathbb{R}^N))$ and that $\varphi \sqrt{\rho_n}$ converges weakly in $L^2(\mathbb{R}^N)$, uniformly in $t \in [0, T]$. Therefore, in order to conclude, we just have to show that we have:

$$\int_{\mathbb{R}^N} \varphi^2 \rho_n(t_n) dx \rightarrow \int_{\mathbb{R}^N} \varphi^2 \rho(\bar{t}) dx$$

whenever $t_n \in [0, T]$, $t_n \rightarrow_n \bar{t}$, and this is straightforward since we have, in view of the above equation:

$$\begin{aligned}\int_{\mathbb{R}^N} \varphi^2 \rho_n(t_n) dx &= \int_{\mathbb{R}^N} \varphi^2(\rho_0)_n dx + \int_0^{t_n} ds \int_{\mathbb{R}^N} \rho_n u_n \cdot \nabla \varphi^2 dx \\ &\longrightarrow_n \int_{\mathbb{R}^N} \varphi^2 \rho_0 dx + \int_0^T ds \int_{\mathbb{R}^N} \rho u \cdot \nabla \varphi^2 dx = \int_{\mathbb{R}^N} \varphi \rho(\bar{t}) dx.\end{aligned}$$

□

References

- [1] D. Bresch, B. Desjardins and C.-K. Lin, On some compressible fluid models: Korteweg, lubrication and shallow water systems. Comm. Partial Differential Equations, 28(3-4) : 843-868, 2003.
- [2] J.W. Cahn, J.E. Hilliard, Free energy of a nonuniform system, I. Interfacial free energy, J. Chem. Phys. 28 (1998) 258-267.
- [3] R. Coiffman, R. Rochberg and G. Weiss, Ann. Math., 103 (1976), pp. 611-635
- [4] F. Coquel, D. Diehl, C. Merkle and C. Rohde, Sharp and diffuse interface methods for phase transition problems in liquid-vapour flows.
- [5] R. Coifman, P.-L. Lions, Y. Meyer, S. Semmes, Compensated-compactness and Hardy spaces, J. Math. Pures Appl., 72 (1993), p 247-286.

- [6] R. Danchin and B. Desjardins, Existence of solutions for compressible fluid models of Korteweg type, *Annales de l'IHP, Analyse non linéaire* 18,97-133 (2001)
- [7] J.E. Dunn and J. Serrin, On the thermomechanics of interstitial working ,*Arch. Rational Mech. Anal.* 88(2) (1985) 95-133.
- [8] E. Feireisl, *Dynamics of Viscous Compressible Fluids-Oxford Lecture Series in Mathematics and its Applications*-26.
- [9] H. Hattori and D.Li, The existence of global solutions to a fluid dynamic model for materials for Korteweg type, *J. Partial Differential Equations* 9(4) (1996) 323-342.
- [10] H. Hattori and D. Li, Global Solutions of a high-dimensional system for Korteweg materials, *J. Math. Anal. Appl.* 198(1) (1996) 84-97.
- [11] David Hoff. Strong convergence to global solutions for multidimensional flows of compressible, viscous fluids with polytropic equations of state and discontinuous initial data. *Arch. Rational Mech. Anal.*, 132(1):1-14, 1995.
- [12] D. Jamet, O. Lebaigue, N. Coutris and J.M. Delhay, The second gradient method for the direct numerical simulation of liquid-vapor flows with phase change. *J. Comput. Phys.* 169(2): 624–651, (2001).
- [13] D.J. Korteweg. Sur la forme que prennent les équations du mouvement des fluides si l'on tient compte des forces capillaires par des variations de densité. *Arch. Néer. Sci. Exactes Sér. II*, 6 :1-24, 1901.
- [14] P.-L. Lions, *Mathematical Topics in Fluid Mechanics, Vol 1*,
- [15] P.-L. Lions, *Mathematical Topics in Fluid Mechanics, Vol 2, Compressible models*, Oxford University Press (1996)
- [16] A. Novotný and I. Straškraba. Introduction to the mathematical theory of compressible flow, *Oxford lecture series in mathematics and its application.* 27
- [17] C. Rohde, On local and non-local Navier-Stokes-Korteweg systems for liquid-vapour phase transitions. Technical report, Math. Institut, Albert-Ludwigs-Universität Freiburg , 2004. Preprint.

- [18] C. Truedell and W. Noll. The nonlinear field theories of mechanics. Springer-Verlag, Berlin, second edition, 1992.